

# **Fisheries, Wetlands and Jobs**

## **The Value of Wetlands to America's Fisheries**

**March 1998**

This report is an update of the 1994 version.

### **Executive Summary**

### **Table of Contents**

Prepared for the Clean Water Network  
by  
William M. Kier Associates  
Consulting Fisheries Scientists  
Sausalito, California

The original 1994 version was written with the assistance of

Clean Water Action  
National Audubon Society  
Natural Resources Defense Council  
National Wildlife Federation  
Sierra Club  
Trout Unlimited  
Campaign to Save California Wetlands

The 1998 version also included the assistance of

Chesapeake Bay Foundation  
Chesapeake Bay Commission

# Executive Summary

*Fishing pumps \$152 billion into the nation's economy and supports about two million jobs.*

The nation is engaged in a continuing debate over the future of America's wetlands - salt marshes, the brackish shallows of bays and estuaries, freshwater swamps, seasonal pools, even Alaska's frozen tundra. The discussion should address the role that wetlands play in groundwater replenishment, flood control, the protection of water quality, the conservation of nature and wildlife - and the production of fish.

The relationships between wetlands and fish production are essential. They are well understood by fisheries experts and most fishermen. This report, prepared from scientific publications and government documents, describes the links between wetlands function and the health of some of the nation's most valuable fish resources.

The values discussed here are substantial. Three quarters of the nation's fish production depends on marshes and other wetland environments. Fishing contributes over \$152 billion to the country's economy and provides jobs for about two million Americans.

*Three quarters of the nation's fish production depends on estuaries, marshes and other wetland environments.*

The U. S. Fish and Wildlife Service estimates that at the birth of our nation, wetlands covered 104 million acres of the contiguous states -- an area roughly the size of California. Half of those wetlands are gone. Although our country is largely developed, we are still destroying wetlands at a rate of more than 100,000 acres year. In a February 1998 Clean Water Action Plan, President Clinton and Vice President Gore vowed to stop the destruction of wetlands in the United States. They announced a multi-agency plan for achieving a net gain of as many as 100,000 acres of wetlands per year by the year 2005. Threatening this goal are nationwide or "quick" permits proposed by the U.S. Army Corps of Engineers, which open the door for increased wetlands destruction.

Wetlands are the critical link in providing clean water for America's families and businesses, as well as supporting the commercial and sport fisheries. The Clinton-Gore Clean Water Action plan has framed the debate over what should be our national wetlands conservation agenda. Congress has an important role to play as well. It must enact strong legislative safeguards to ensure long-term protection for wetlands. This report makes clear the essential role wetlands must play in a sustainable American sport and commercial fishing industry.

[Table of Contents](#)

[Turn the page...](#)

# Fisheries, Wetlands, and Jobs

## Table of Contents

[Executive Summary](#)  
[Introduction](#)  
[Land of Pollock and the 1% Rule](#)  
[Restoring the King](#)  
[Dungeness Crab](#)  
[California's Halibut](#)  
[Gulf Shrimp](#)  
[Menhaden: A Little Pogy in All of Us](#)  
[Trout Fishing Michigan's Wetlands](#)  
[Dollars and Sense on the Big Muddy](#)  
[Trouble in Paradise](#)  
[Leapin' Bluefish](#)  
[Stripers on the Chesapeake](#)  
[Lobsters: Home for Homarus](#)  
[Conclusions](#)  
[About This Report](#)  
[References](#)

# Introduction

This report explains the relationship that fish have with their wetland habitats and the relationship that we have with fish. Both relationships are important to survival - theirs and ours.

Fish, like many humans, may change the location and nature of their habitat as they go through stages of development. For example, many ocean fish spend their youth in shallow-water wetland habitats and can venture into the open sea only when they have become strong swimmers. Scientists believe, and have in many cases clearly demonstrated, that the availability of habitat and the survival of fish are absolutely linked. The disappearance of wetlands, therefore, leads to the decline of the fish that depend upon them.

*We sold \$8.7 billion of our fish products to foreign trading partners; about \$6 billion of them were produced in the wetlands.*

Wetlands are an essential habitat for both salt and freshwater fish. While we may struggle for a politically acceptable definition of wetlands, their nature and function are very clear where fish are concerned. Wetlands cycle nutrients out of mud, sand and water back into bays, lakes and streams, promoting the growth of the smallest organisms in the aquatic food chain. These small creatures are eaten by larger ones and then by fish. Such "primary production" and predation work especially well where the water is sheltered, shallow and rich with nutrients. Therefore, no matter how we may define them, wetlands are vital to fish for food and shelter.

The term "fishery" puts us in the picture. It refers to our harvest of fish resources, for food or sport. We talk of an "oyster fishery", a "catch-and-release trout fishery", or the "Columbia River salmon gillnet fishery" as the place of the harvest and even the harvesting method.

Fishing is big business in this country. Our commercial fishermen harvested nearly ten billion pounds of fish in 1996. National Marine Fisheries Service scientists estimate that nearly seven billion pounds of these fish depend upon inshore-wetland habitats.

One hundred thousand U.S. fishing boats are the workplace for a half million fishermen and fisherwomen. Most of these boats are family businesses, similar to family farms or other small businesses ashore. Each of the onboard jobs supports several workers on land, including those who unload, process and truck fish, and those who fuel, provision and repair the boats.

*Sports fishermen's \$37.8 billion in spending means \$108.5 billion in economic output, 1,210,083 jobs and \$5.4 billion in state and federal taxes.*

The U.S. Department of Commerce reports that Americans spent \$41.2 billion on seafood products in 1996, a six percent increase over 1995. \$34.4 billion was spent on domestic seafood and \$6.8 billion was spent on imported seafood. We also sold \$8.7 billion worth of fishery products to foreign trading partners, about \$6 billion of which were produced in wetlands.

But that doesn't tell the full story. Available national data on the commercial fishing industry from the U.S. Department of Commerce does not report the full economic output of this industry. The \$43.1

billion value assigned to the domestic commercial fishing industry underestimates the total economic value of that industry because expenses such as boat repairs, boat purchases, fuel, etc., have economic multiplying effects that have not been included in their analysis. This is because economic multipliers for the commercial fishing industry vary by fishery and by region, and there is no single multiplier for this industry. Some states have done economic analyses with multipliers, but we have not uncovered any national data of this type for the commercial fishing industry. Therefore, the total economic output of this industry is really much higher than reported.

[Table of Contents](#)

[Turn the page..](#)

# Land of Pollock and the 1% Rule

Alaska tops the nation with the extent of its wetlands and the rate at which it is destroying them. Development has destroyed more than 50 percent of the historic wetlands surrounding Juneau and Anchorage. The destruction of Alaska's wetlands places the state's three billion dollar a year fishing industry at risk.

*The destruction of Alaska's wetlands places the state's three billion dollar a year seafood industry and 55,000 jobs at risk.*

The loss of Alaska's wetlands may seem justifiable considering that half the state, at least 170 million acres, is covered with them. Nearly all these acres are inland wetlands, bogs, muskegs, tundra, swamp and forests essential for wildlife and fish such as salmon. The state has only 345,000 acres of salt marsh wetlands. This is where politics and ocean fish production come into conflict.

Alaska promotes a "one percent rule." The rule allows that Alaska be held to a different wetlands protection standard than the lower 48 states, until the state has eliminated one percent of its historic wetlands. Alaska's one percent wetlands exemption was adopted by the Bush Administration in November, 1992.

Alaska's population has tripled since 1950. Most of this growth adjoins coastal waters. Most of the development pressure on Alaska's wetlands is, therefore, on its coastal wetlands.

Simple arithmetic clearly demonstrates the problem. Alaska's 345,000 acres of coastal wetlands represent only two-tenths of one percent of the state's total. The "one percent rule" would allow the destruction of five times the total area of Alaska's coastal wetlands. The "one percent rule" draws no distinction between the more plentiful inland wetlands and the far less plentiful – and more critical to fishing – coastal wetlands.

Fishing is serious business in Alaska. The state's Department of Commerce and Economic Development reports that more than one in ten Alaskans makes his living from the fisheries. In addition, ships and workers from many other states and nations make seasonal journeys to Alaska to participate in the harvest. Included in this work force are the crews of nearly 18,000 vessels which land six billion pounds of seafood a year, 60 percent of the nation's total, plus workers at more than 400 fish handling and processing plants.

While Alaskans debate the future of their coastal ecosystems, evidence linking wetlands to the production of the state's most valuable fish grows. The results of research by state and federal fisheries agencies shows that Alaska's salmon, steelhead, herring, sole, and flounder are all strongly dependent on wetland habitats.

According to a 1991 study by the National Marine Fisheries Service, 76 percent of Alaska's seafood harvest came from inshore and wetlands-dependent species in that year. While more recent numbers are not available, the importance of this relationship remains clear.

Alaska's fastest growing ocean fishery is for walleye pollock, a member of the cod family. According to the National Marine Fisheries Service (NMFS) two and a half billion pounds of Alaska pollock were landed in 1996, a catch worth a quarter of a billion dollars.

Scientists from the government's Auke Bay Laboratory near Juneau regularly survey the region's shallow waters. Young pollock and ocean perch, another economically vital species, are consistently found in the shallow waters around Juneau and other coastal population centers. These shallow water habitats will be degraded unless their value to Alaska's fisheries is recognized in the wetlands regulatory process.

[Table of Contents](#)

[Turn the page...](#)

# Restoring the King

Redwood Creek nestles in the coastal rain forest of northwestern California halfway between Humboldt Bay and the Oregon border. The stream was once famous for its fishing. Redwood Creek yielded chinook salmon - also known as kings because of their great size - weighing up to 65 pounds, in addition to coho salmon and steelhead.

*Studies link the salmon decline to the loss of wetlands.*

All three salmon runs have now declined dramatically. The chinooks - the kings - have been hit the hardest. Their numbers have plummeted 80 percent since the late 1960s.

The land around Redwood Creek has been acquired by the government for the Redwood National Park. National Park Service scientists began to study Redwood Creek more than 20 years ago to learn what could be done to return the kings to the stream. These studies soon determined that the disappearance of salmon from Redwood Creek was linked to the loss of wetlands from the stream's lower reaches.

Floods swept through lower Redwood Creek in the winter of 1965, closing the Redwood Highway and damaging the streamside village of Orick. Congress reacted swiftly, authorizing the Corps of Engineers to straighten and levee Redwood Creek in order to speed its flow directly into the Pacific Ocean.

Before the Corps' 1968 channelization project, lower Redwood Creek spread into side channels and backwaters. It was in these wetland areas, Park Service biologists concluded, that young chinook salmon lingered on their migration to the sea. This was where the fish had fed and gained the strength to survive in the open ocean.

The Park Service settled on a plan more than fifteen years ago to improve Redwood Creek's fish habitat. The Service estimated it will take at least \$2 million to modify the now 30-year-old flood control project to restore some of the lost wetland rearing areas. Fishery experts agree this is the only way to return the kings to the stream.

In stark contrast to the speed with which the Corps of Engineers' levee project was funded, the Park Service's request for funds to restore Redwood Creek's wetlands has moved at a snail's pace. In terms of the National Park system's nationwide priorities, it seems the return of the king to Redwood Creek must wait.

There are hundreds of streams like Redwood Creek up and down the Pacific Coast where habitat for young salmon must be pieced back together if salmon are to continue to contribute to the nation's economy. Nearly \$100 million was spent to repair these battered streams in California alone over the past 20 years. Much of this money was raised by a tax on the state's commercial salmon fishermen's own landings, even though both fish and work have declined steadily since 1978.

The fishermen are fighting for their lives. The industry had gained a major legislative victory in 1992 when it persuaded Congress to restore a significant amount of water from the vast federal Central Valley Project back into key California rivers for salmon restoration. The biggest program under consideration now in California for wetlands and fishery restoration is known as CalFed. This program has a potential one billion dollars earmarked from state and federal coffers for restoring wetland and river habitat.

The rationale supporting these efforts is strong. After analyzing the state's salmon problem, the California legislature concluded that rebuilding salmon runs to twice the depressed 1980s levels would provide economic benefits of \$150 million a year.

Full implementation of the doubling effort over several years would yield six billion dollars in net profits to the state, one billion dollars of which would go to small businesses. In addition, it was found that the salmon restoration objective, which Congress incorporated into the 1992 Central Valley Project reform, would create 8,000 new jobs.

The bright promise of the Central Valley Project Improvement Act (CVPIA) seems to be slipping away. The Administration's CVPIA policy commits to using only about half of the freshwater flows dictated by the law for salmon, and is now the subject of lawsuits filed by environmental and fishing interests. The risk with CalFed is that money will be spent on restoration projects without providing the key element for wetlands and river habitat: water.

Even if the money for restoration becomes available, this should not become the excuse for destroying wetlands further. In a 1992 report to Congress, the National Research Council's Committee on Restoration of Aquatic Ecosystems cautioned, "[M]any [restoration] projects fall short of the goal of returning ecosystems to the predisturbance condition, and there is indeed considerable controversy over whether or not wetlands can actually be restored [at all]." As Redwood Creek demonstrates so poignantly, it is a whole lot easier to protect a wetland than it is to put one back together.

# Dungeness Crab

Dungeness crab are near the top of life's blessings. Sweet and succulent, Dungeness crab meat, like fresh ocean-caught king salmon, is one of the nation's finest seafoods.

When northern California's rivers swell with late rains and snow-melt, the strength of their outflow forces incoming tides to the bottom of bays and estuaries. These bottom tides, or gravitational currents, carry the crab larvae from the ocean through the Golden Gate and deposit them among the wetlands of the upper San Francisco Bay system.

These springtime travelers look more like tiny shrimp at this megalops stage. The wetlands nourish the young crabs which grow, molt (shed their shells) and grow some more until they reach adulthood in the Bay a year or so later. To be precise, not all Dungeness crabs rear in the Bay. Some appear never to leave their nearshore ocean spawning grounds. Those that do rear along the edges of San Francisco Bay's sheltering wetlands grow more quickly and doubtless survive at a higher rate, contributing more to the fishery.

Young Dungeness crabs do not appear to use areas of the estuary where salinities are less than about one-third that of seawater. To the extent young crabs depend on wetlands, and they certainly appear to, they require areas bordering salty, rather than fresh or brackish, water. Therefore, it is not possible to mitigate the destruction of Dungeness crab-sustaining wetlands at sites that are less than one-third seawater.

California fishers landed 12.3 million pounds of Dungeness crab in 1996, up 35 percent from 1995 and worth more than \$16 million dockside. A healthy Dungeness crab fishery provides some relief to those fishermen who have traditionally depended on salmon for a living.

[Table of Contents](#)

[Turn the page...](#)

# California's Halibut

*Commercial catches of halibut have declined from nearly 5 million pounds to just over one half million pounds a year since 1920.*

Westerners have a hearty appreciation for their seafood, including halibut. While Pacific halibut from Alaska is excellent, fresh California halibut is especially prized. These flavorful fish, which grow to more than 50 pounds, range from northern Washington to Baja California. Much of the California halibut catch now occurs, however, inside the Southern California Bight, the nearshore waters between Point Arguello and Mexico.

Like many of California's fisheries the halibut industry was launched from San Francisco. Italian-born fishermen began towing a bottom net for halibut, a paranzella, between two lanteen-rigged sailboats during the 1880s. The fishery soon grew north to Bodega Bay and Eureka and south to Santa Barbara, Los Angeles and San Diego. By 1920 California halibut landings had swelled to nearly five million pounds a year.

Declining numbers of halibut and the resulting protective regulations have taken their toll on the halibut fishery. Commercial catches have declined in recent decades to just over a half million pounds a year. In 1994, commercial fishermen took 510 thousand pounds of halibut worth an estimated \$1.3 million. The fish continues to be a popular target for sports fishermen. The loss of wetlands to development along California's coast has contributed significantly to the decline in halibut.

Like many ocean fish, mature halibut broadcast their eggs in shallow waters where they become part of the planktonic mix, to be tossed about by winds and currents. Researchers have begun to pinpoint when and where larval halibut settle out along the coast to begin their growth needed to survive to harvestable size. The new information make clear the importance of wetlands to this fishery.

Researchers from the State University system's Southern California Ocean Studies Consortium have captured young halibut in a variety of open water and protected areas along the Bight during the 1980s. The samples from these surveys makes clear the high preference of young halibut for protected shallow water areas such as those provided by salt marshes. The loss of these wetlands - 90 percent of the bays and estuaries in southern California have been severely altered or destroyed by human activities - is clearly implicated in the decline of California halibut in recent decades.

The small fraction of historic wetlands that remains along the southern California coast will have to be protected vigorously if California halibut are to continue to grace the tables of our nation.

[Table of Contents](#)

[Turn the page...](#)

# Gulf Shrimp

About a third of a billion pounds of shrimp are caught by shrimpers across the nation each year. At \$1.50 to two dollars a pound dockside, the nation's annual shrimp harvest puts a half billion dollars in the pockets of U.S. fishermen. The nation's most productive shrimp fishery is the Gulf of Mexico where fishermen from Texas, Louisiana, Mississippi, Alabama and Florida harvested 438 million pounds of brown, white and pink shrimp in 1995 and 1996 combined, worth over \$838 million dockside according to the U.S. Department of Commerce.

*The nation's annual shrimp harvest puts a half billion dollars in the pockets of U.S. fishermen.*

The dockside value of Louisiana's commercial shrimp harvest only tells a small part of the story of this fishery's value. For example, in one small fishing community, Lafitte - Barataria, roughly twenty families operate shrimp boats in the Gulf which are each worth a half million dollars. About 250 families in the community own smaller skiffs for fishing inland waters, ranging in value from a thousand to one hundred thousand dollars apiece. A comprehensive estimate of the importance of commercial fishing to the state's economy would figure in these boats along with boat building and repair shops, the fuel suppliers and the many other parts of the local economy that depend on commercial fishing.

Nowhere in the nation is the link between wetland habitat and fish production more obvious than in the Gulf, where National Marine Fisheries Service scientists estimate that 98 percent of the harvest comes from inshore, wetlands dependent fish and shellfish. Gulf shrimp clearly head the list of the region's wetland dependent species.

The mature shrimp spawn in the Gulf's offshore waters. Fertile eggs soon hatch into free-swimming larvae, and the larvae quickly pass through a series of molts. During the postlarvae stage the shrimp enter the estuaries along the coast to become bottom feeders.

In the estuaries the juvenile shrimp feed at the marsh-water or mangrove-water interface or in submerged seagrass beds. These areas offer a concentrated food supply of detritus, algae and microfauna and some protection from predators. Both the growth and survival of the young shrimp are largely dependent on local salinity and temperature regimes.

In its shrimp management plan the federal Gulf of Mexico Fishery Management Council notes that, "The weakest link in the life cycle chain is the estuarine phase of growth." Man-caused factors implicated by the Council in the decline of Gulf shrimp habitat were "bulkheading that removes critical marsh-water and mangrove-water interfaces" together with "alterations in freshwater discharge that create unfavorable salinity regimes."

The Everglades of south Florida, Marjory Stoneman Douglas' River of Grass, have for centuries fed freshwater through the mangroves into the warm tides of the Gulf and Florida Bay. The seagrass beds of this region are critical habitat for pink shrimp. The diversion of Everglades freshwater to urban development and as a result of drainage programs has driven up salinities in the receiving bays, destroying the seagrass shrimp nurseries. By 1990, 10,000 acres of Florida Bay seagrass beds had been lost completely while another 50,000 acres was in serious decline. Recent research shows that while lost seagrass acreage has recovered in some areas, other areas have continued to decline, indicating an overall net loss that has yet to be fully quantified.

Because the wetlands-shrimp production relationship is so strong it illustrates an important matter in the debate over wetlands: the boundaries of wetlands are not always simple to define. Freshwater from the heart of the Everglades' huge wetlands system moderates the salinity of Florida Bay more than 50 miles downstream. Moderate salinities are needed for healthy seagrass beds. Agriculture and urban development rob the Everglades of much of this water, and the water that returns is often polluted. As seagrass beds die off due to salinity fluctuations and nutrient pollution, the sediment held by their roots is suspended, blocking out the sunlight needed by the remaining plants – a downward spiral.

Florida Bay's seagrass beds are prime pink shrimp habitat. The Everglades, the Florida Bay seagrasses and the Gulf's offshore shrimping grounds are, in this way, an interdependent ecosystem – all parts of which need protection.

Many of the 14,000 jobs in more than a thousand Gulf Coast fish processing plants are sustained by the shrimp fishery. And commercial fishing represents only a part of the economic engine fueled by these tasty invertebrates. Shrimp are prime prey for the area's gamefish and, therefore, popular as bait with the region's anglers.

According to the Census Bureau, Gulf state sport anglers brought \$7.9 billion to the region's economy in 1996. Florida anglers led the way, with nearly \$3.3 billion dollars in total expenditures. Anglers in Texas (\$2.9 billion), Louisiana (\$824 million), and Alabama (\$835 million) also provided a significant boost to the regional economy.

[Table of Contents](#)

[Turn the page...](#)

# Menhaden: A Little Pogy in All of Us

Until the recent boom in Alaska pollock, menhaden -- "pogy" to fishermen -- made up an impressive one-third of our nation's industrial catch. Because we don't eat menhaden, or we don't know that we do, the fish attracts little interest outside a few southern coastal towns where malodorous menhaden plants exude the "smell of money."

The menhaden fishery began in New England. Colonists used the fish for fertilizer and bait. Rhode Islanders began pressing lamp oil from menhaden about 1800. Beginning in the 1930s the bulk of the fishery shifted to the South, where it now involves two species, Atlantic and Gulf menhaden.

Menhaden are cooked and pressed into meal, oil and "solubles." Mixed into poultry feeds, the meal and oil are vital for improving growth rates and food efficiency. The solubles are blended with soybean meal for cattle and swine feed. Menhaden oil is mixed with other fats in cooking oil, shortening and margarine, marine lubricants, plasticizers, alkyd resins, oil for paint and even lipstick.

Increasingly mechanized, the fishery employs more than 100 boats, 1,400 fishermen and a like number of plant workers. The 1.8 billion pound 1996 menhaden harvest brought fishermen \$94 million and guaranteed products essential to the South's poultry producers and many other industries.

Menhaden absolutely need wetlands to exist. Tragically, their coastal nurseries are under siege. Researchers estimate that Louisiana's coastal marshes, critical habitat for menhaden, are being lost to a combination of natural and development forces at an average rate of between 20,000 and 25,000 acres a year.

In recent years, an alarming new threat has emerged. A mysterious micro-organism, the dinoflagellate *Pfiesteria piscicida* -- the "cell from hell" -- has caused large kills of menhaden and other fish, and has even been associated with human illness. Researchers believe that this organism may have always been present in estuaries in a dormant or non-toxic state.

The concern now is that nutrient pollution of rivers and estuaries from agriculture, livestock and urban development may have caused the broad scale emergence of the deadly form. Restoring wetlands, which help trap nutrients, is one way to cut down on polluted runoff going into the estuaries where menhaden are found.

[Table of Contents](#)

[Turn the page...](#)

# Trout Fishing Michigan's Wetlands

Fishing is big business in Michigan. Angling in Michigan contributed \$2.8 billion to the economy in 1996 and generated about \$200 million in state and federal taxes.

*Angling contributed \$2.8 billion to Michigan's economy in 1996 and generated about \$200 million in taxes.*

Michigan owes some of its best fishing to wetlands that would be stripped of federal protection under an ongoing series of proposed policy changes to the U.S. Army Corps of Engineers nationwide permit system, which provides specific provisions for protecting wetlands. Fortunately to date these proposals have not been accepted. A look at Michigan's Pere Marquette River will explain how wetlands serve freshwater ecosystems, and how proposed changes to federal wetlands protection regulations threaten these resources.

The Pere Marquette River rises from two large wetland complexes, the Oxford swamp and the Widewaters. Along its 80 mile trip to Lake Michigan, the river takes on water from a number of smaller wetlands. At some points the entire river must work its way through a mixture of swamps, floating bogs and marshes.

The water chemistry of the Pere Marquette rarely varies. The river never freezes and it rarely gets above 60 degrees in the summer. The wetlands release the water they store up in winter and spring gently throughout the year. In short, the Pere Marquette is prime trout water and its brown trout and steelhead fishing attract fishermen by the thousands from all over the country.

Strong, enforceable wetlands protections are critical to the water sources that make the Pere Marquette, and hundreds of other wetland-origin trout streams in the Midwest and Rocky Mountain regions of the nation, so valuable. Although the Pere Marquette's wetlands have saturated soils, the water seldom is at or above the surface. The Corps of Engineers has recently proposed new nationwide permits that would make it easier to fill the wetlands that feed the Pere Marquette and to disrupt the flow of water from those wetlands to the river. These nationwide permits cover a variety of activities in the wetlands and along the river, such as residential and commercial development, road building, oil and gas drilling, and gravel mining. Developers and industries engaging in these activities could fill significant amounts of wetlands without the need to apply for individual permits, and the Corps could give such projects the go-ahead without the full public notice required by the individual permit procedure.

The Corps is scheduled to release this proposal in March for public comment. People who are concerned about preserving habitat for trout should keep a close eye on this proposal.

As the nation considers how to cope with the damage from the devastating floods exacerbated by El Nino, we should keep in mind an essential function of wetlands. They soak up water like giant sponges when it might otherwise race away in a flood, and they release that water slowly, benefiting man and beast. That is the way the Pere Marquette works, and that is the way the floodplains of the Missouri River and other Midwest streams may be made to work once more.

The President's Clean Water Action Plan, announced in February 1998, sets an ambitious goal of creating a net gain of 100,000 acres per year – by reducing losses and accelerating restoration. The plan is a dramatic step in the right direction that deserves the support of Congress.

[Table of Contents](#)

[Turn the page...](#)

# Dollars and Sense on the Big Muddy

*The Flood of '93 cost taxpayers nearly \$16 billion in disaster recovery payments, much of which was lost in 1995 when floods again dismantled levees.*

Throughout time Missouri River fish have danced to the seasons. Bass, catfish, sturgeon and freshwater drum all thrived as the "Big Muddy" swelled across its floodplain, gathered up organic decay and nutrient-rich silts, gorged underground aquifers and saturated its wetlands. These flooded backwaters, teeming with young fish, then drained slowly, relentlessly back to the main river, feeding its life web with water, salts and silt.

For more than a century now men have worked to herd the Missouri between levees, drain its wetlands, and strip its riparian forests for farmland. Ninety percent of the Missouri's historic bottomlands have been obliterated. The river's commercial fish harvests have plummeted 80 percent in the past 50 years. Rather than support a potentially lucrative caviar fishery, the pallid sturgeon has been listed as "endangered" under the federal Endangered Species Act.

When the Missouri and Mississippi rivers overwhelmed their flood control works in 1993 the nation's taxpayers got a cleanup bill for \$16 billion. When these rivers turned right around and did the same thing two years later, it seemed like it was time to rethink the way we were spending money.

In 1997 the U.S. Fish and Wildlife Service proposed that 45,000 acres of historic Missouri River floodplain be rejoined to the river by expanding the existing 14,000-acre Big Muddy National Fish and Wildlife Refuge. The decision is still pending, but it is the preferred alternative included in the environmental impact statement for the Big Muddy. The lands will be acquired from willing sellers weary of battling the river. The refuge expansion idea looks like a winner from at least two standpoints: Missouri's own fishing and the health of the larger Mississippi River ecosystem.

Missouri is a fishing state. Spending by anglers gave the state an economic boost of \$1.5 billion in 1996. The Big Muddy National Fish and Wildlife Refuge expansion will increase the river's fish production and thereby expand both angling and commercial fishing opportunities.

American taxpayers are eyeing their bills a lot more closely these days. The benefits of the Big Muddy wetlands restoration will go far beyond Missouri and, unlike the flood control works they replaced, they will endure.

[Table of Contents](#)

[Turn the page...](#)

# Trouble in Paradise

Louisiana's vehicle license plate proclaims that the state is the "Sportsman's Paradise". The numbers would certainly suggest so. The state is blessed with 42 percent of the nation's coastal wetlands. These include the central Gulf of Mexico area known as the "Fertile Crescent", some of the most productive fisheries habitat in the world. Louisiana leads the lower 48 states in total fisheries production and in the production of shrimp, blue crabs and oysters.

*Sports anglers in Louisiana spent \$824 million in 1996. The state's commercial fisheries generated \$2.1 billion in retail sales that year.*

Louisiana sports anglers spent \$824 million in 1996 resulting in a total economic output of \$1.54 billion to the economy. The state's commercial fishing industry had retail sales of \$2.1 billion in 1996 with a total economic effect of \$2.8 billion according to the State of Louisiana Department of Wildlife and Fisheries.

Louisiana is losing coastal wetlands, however, at the rate of up to 35 square miles a year. That rate has more than doubled since the late 1940s, when work on the Mississippi River flood control projects accelerated. The project levees cut Delta wetlands off from the river's sediment supply, and it was those very sediments that helped build the river's Delta wetlands over the ages.

As Louisiana's wetlands go, so goes southwestern Mississippi. The two states share the region's wetlands blessings - and the tragedy of their loss. The region has campaigned successfully for federal recognition of the problem. Congress passed the Coastal Wetlands Planning, Protection and Restoration Act - the "Breaux Act" - in 1990. The act calls for the implementation of specific projects to improve the delivery of river-borne sediments into the threatened coastal wetland areas. The main projects called for in the act have proven controversial, however, and progress since 1990 has been limited to minor measures.

Noting the then-incipient shrinkage of Louisiana's coastal wetlands back in 1928, one researcher urged "a broad program of conservation along new and intelligent lines", concluding that wetlands loss "should be considered a state and national problem ... to the end that the state and nation may enjoy a more balanced diet, more healthful recreation, and enduring prosperity." Those words ring every bit as true today as they did 70 years ago. Time is running out for Louisiana's coastal wetlands and her enormously valuable wetlands-dependent fisheries. We can only pray that the region's political leadership will grasp those "new and intelligent lines" before Paradise is lost.

[Table of Contents](#)

[Turn the page...](#)

# Leapin' Bluefish

*The destruction of inshore habitat, including the pollution of bays and estuaries, is strongly implicated in the blues' decline.*

The scientific name for bluefish, *Pomatomus saltatrix*, salutes the fish's impressive energy. While the first name attempts to describe the dark, scar-like line on the gill cover ("cover-cut"), *saltatrix* exclaims "somersaulter" - leaper. The ubiquitous blue bites hard and fights valiantly, making him a favorite of saltwater anglers from the South Atlantic coast to New England. Unfortunately, blues are in decline and deteriorating shoreline conditions may be the reason.

Adult bluefish move around the ocean. They winter off the coast of Florida and head north in the spring. A major bluefish spawn occurs in early spring just landward of the Gulf Stream between Florida and southern North Carolina. A second summer spawn takes place off the mid-Atlantic. Larvae from the spring spawn head north with the currents, around Cape Hatteras to the offshore waters of the mid-Atlantic Bight. As the waters nearer the shore warm in late spring, juvenile blues move into bays and estuaries to feed on wetlands invertebrates such as opossum shrimp, amphipods and small wetlands fish like silversides and killifish.

Bluefish are famous for feeding frenzies in which they attack schools of baitfish and tear their prey savagely to bits. The scraps attract gulls, which fishermen seeking blues eagerly track, seeking a chance to join the fray.

Recreational anglers caught 93 million pounds of bluefish in 1983. By 1993, the catch plummeted to 20.6 million pounds. "We have strong reason to believe the population is declining" State of Connecticut fisheries research supervisor Victor Crecco told the press that year.

The evidence behind Crecco's assertion continues to mount. Bluefish catches have continued to decline, falling to just over 15 million pounds in 1996. While heavy fishing pressure certainly shares some responsibility of reduced catches, researchers believe that the destruction of habitat and pollution contribute to the problem.

Despite these hard times, blue fever remains strong. More than 60 percent of bluefish caught in 1996 were taken from party or private vessels, an important source of employment for coastal communities on the eastern seaboard.

[Table of Contents](#)

[Turn the page...](#)

# Stripers on the Chesapeake Bay

*Wetlands within the Bay's  
64,000 square mile  
watershed must be protected  
vigorously.*

There are three primary striped bass populations along the Atlantic coast: Hudson River, Chesapeake Bay and the Roanoke River. For generations striped bass, also known as rockfish, sustained major commercial and recreational fisheries. The popularity of stripers with anglers often drove annual sport catches beyond those of the commercial fishermen. The commercial striped bass harvest peaked in 1973 at about 14 million pounds, and then began a precipitous decline.

Following years of particularly poor survival of striped bass, the result of poor water quality and severe fishing pressure, Congress passed the Striped Bass Conservation Law in 1984. This legislation directed the Departments of Commerce and Interior to assist the interstate Atlantic States Marine Fisheries Commission to moderate fishing pressure in order to restore striper populations to healthy levels.

Although prudent management appears to be making a positive contribution, fishing pressure is not the only problem. Most striped bass spend much of their life cycle in coastal waters, migrating into tidal freshwater tributaries to spawn. Healthy spawning habitat and other environmental conditions are critical to the reproductive success of striped bass.

Water pollution has been especially hard on the Chesapeake Bay stripers. The overall total load of nutrients received from all sources flowing into the Chesapeake Bay adds up to about 25.8 million pounds of phosphorus and 362.4 million pounds of nitrogen a year according to U.S. Environmental Protection Agency 1998 estimates. These nutrients stimulate the growth of algae near the water surface, blocking sunlight needed by the bay's underwater sea grasses – critical rearing areas for juvenile stripers. When the mats of algae die they consume the oxygen needed by seagrass and stripers alike. By the late 1980's, half the bay's seagrass beds had been lost in this way.

The magnitude of the nutrient problem prompted the inclusion of a specific commitment in the 1987 Chesapeake Bay Agreement to reduce by 40 percent the amount of "controllable" or "man-made" nitrogen and phosphorus entering the bay by the year 2000. A 1997 review of the Chesapeake Bay nutrient problem conducted by the U.S. Environmental Protection Agency indicated that the nitrogen loading goal would not be met without substantial effort. While the review concluded that the phosphorous loading goal would be met by the year 2000, it would begin to rise again if nothing changes.

The confirmed presence of the toxic micro-organism *Pfiesteria piscicida* after a fish kill on Maryland's Pocomoke River last summer reinforces concern about nutrient levels in the Bay. Many fishermen and conservationists blame the watershed's large number of chicken farms for sending slugs of excess nutrients into the Pocomoke, especially during the unusually heavy rains of 1996. They see the outbreak of this organism and the fish kill as a result of that pollution. Quoted by the press when the story broke, Mike Hirshfield, vice president for Chesapeake Bay Foundation, the region's largest environmental group, said, "As if we needed another reason to worry about animal waste nutrients, this was a real kick in the teeth."

Time has shown that reducing the striper harvest until the stock could recover, combined with efforts to reduce pollution and improve estuarine habitat has resulted in a remarkable comeback for striped bass. The stock was declared restored as of 1995. A modest recreational and commercial fishery for stripers has been reopened, and the fishery is being monitored closely to assure that it is kept at sustainable levels.

The wetlands within the Bay's 64,000 square mile watershed must also be protected vigorously for their nutrient trapping and removal capabilities. While the net loss of estuarine wetlands has been slowed, the loss of freshwater wetlands in the Bay watershed is accelerating -- a cause for concern and continued vigilance.

[Table of Contents](#)

[Turn the page...](#)

# Lobsters: Home for Homarus

The American lobster, *Homarus americanus*, plays an important role in the culture and heritage of New England. This fishery is also important to the regional economy. Each year, the fishermen of Massachusetts, Maine and Connecticut harvest this succulent delicacy by the millions. Yearly catches in this decade have averaged better than 60 million pounds.

*In 1996, New England fishermen landed 71.6 million pounds of lobster, worth nearly \$242 million dollars.*

The lobster is not usually thought of as a wetlands-dependent animal. And that, according to New England researchers, is an oversight that could undermine the region's valuable lobster fishery. Inshore areas that have shallow water, lowered salinities, varying temperatures and increased turbidity may be precisely what are needed to speed young lobsters on their way to becoming "keepers."

Recent research shows that juvenile lobsters inhabit inshore areas including cobble, eelgrass and salt marsh peat reefs in relatively high densities. There is strong evidence that years of abundant freshwater flow from wetland areas into these inshore habitats are followed nine years later -- the period it takes juvenile lobsters to grow to legal size -- by good lobster harvests.

Better survival of juveniles to harvestable size, "recruitment" to fisheries managers, is essential if lobster fishing is to be sustained. The fishery is currently taking a relatively high percentage of the adult lobsters, the breeding population needed to sustain future harvests. Fisheries management will play a pivotal role in the future health of these resources, even as the connection between wetland functions and lobster production is coming under closer scrutiny. Pending the scientists' final verdict, lobstermen and their friends will do well to continue their support for wetland conservation efforts.

[Table of Contents](#)

[Turn the page...](#)

# Conclusions

From hundreds of fishing communities across the land and thousands of coastal and river fish habitats, the evidence mounts steadily that most of this country's \$152 billion dollar fisheries economy - and the approximately two million jobs that go with it - depend on wetlands.

Where wetlands have been destroyed, fisheries have declined. Some heroic efforts, like those of California's salmon fishermen, are being made to restore wetland and fish resources. In relatively undisturbed areas like Alaska, the wetlands-fisheries connection appears to be inadequately appreciated. In the New England lobster fishery the importance of wetlands, always a fact, is only now being recognized.

The fishermen of this country are literally fighting for their lives. Those with organizing skills and some remaining resources are leading -- even winning -- conservation battles, as in the case of the recent reform of California's federal Central Valley Project. In too many cases, however, fishermen's pleas for wetlands protection have been lost in the nation's race to develop coasts, estuaries and rivers for agricultural, energy, recreational and urban projects.

America's fisheries are sustainable, but only if the habitat they need is identified and protected. Fishery declines will continue unless the rebuilding process tackles habitat problems as well pollution and overfishing.

The alarming outbreak of *Pfiesteria* in the estuaries of the Atlantic and the Gulf coasts drives home the point that taking our wetlands for granted threatens our economic, cultural and physical health.

The President's Clean Water Action Plan sets an ambitious goal of creating a net gain of 100,000 acres of wetlands a year by the year 2005 -- by reducing losses and accelerating restoration. The plan is a dramatic step in the right direction that deserves the support of Congress.

However, in contradiction of the President's goal, the U.S. Army Corps of Engineers recently drafted new proposed nationwide "quick" permits that would make it easier to destroy wetlands. These nationwide permits cover a variety of activities in wetlands such as residential and commercial development, road building, oil and gas drilling, and gravel mining. Developers and industries engaging in these activities could fill significant amounts of wetlands without the need to apply for individual permits, and the Corps could give such projects the go-ahead without the full public notice required by the individual permit procedure.

The Corps is scheduled to finalize this proposal in March 1998, after which it must provide the proposal for public review and comment before making a final decision by the end of 1998. People who care about wetlands should follow this proposal closely and make their voices heard to ensure that these permits do not foster greater wetlands destruction with resulting harm to fisheries and the jobs that depend on them.

As the nation considers how to cope with the damage from the devastating El Nino, we should keep in mind an essential function of wetlands. They soak up water like giant sponges when it might otherwise race away in a flood, and they release that water slowly, benefiting people, fish and wildlife.

Wetlands are absolutely essential to three-quarters of America's fishery production. Wetlands are complex ecosystems that cycle water, air, soil, nutrients and energy into the waters and life that surround them. It is not surprising that the National Research Council questions whether we can create or restore wetland ecosystems. The prudent course, clearly, is to protect our remaining natural wetlands.

[Table of Contents](#)

[Turn the page...](#)

# About this Report

This report was originally prepared by William M. Kier in 1994 for the Campaign to Save California Wetlands and was funded by a grant from the David and Lucille Packard Foundation. The updated version of this report was prepared by Eric G. Doyle with Mr. Kier's supervision, for the Clean Water Network. It was paid for by general support grants from the Pew Charitable Trusts, the Moriah Fund, C.S. Mott Foundation, the Town Creek Foundation, and the Homeland Foundation.

Mr. Kier is a certified fisheries scientist with more than 35 years of professional experience in the protection and management of freshwater, estuarine and marine sport and food fish resources. His early work in the field included life history studies of the fishes of the San Francisco Bay-Delta Estuary. He became assistant chief of the California Department of Fish and Game's environmental protection branch, the state's assistant Resources Secretary and the principal fisheries advisor and director of research for California's State Senate. The consulting firm he leads specializes in fisheries management analysis and restoration planning.

Eric Doyle has a B.S. in marine biology, and a Masters degree from the School of Marine Affairs - University of Washington, specializing in the economic and policy implications of natural resource management issues. Mr. Doyle has worked extensively with the National Marine Fisheries Service on Endangered Species Act listings of salmon and steelhead in the Pacific Northwest and California.

The Clean Water Network is comprised of more than 1,000 local, state, regional and national groups nationwide working to strengthen federal clean water policies so that our nation's waters will be fishable and swimmable. Member groups include a variety of organizations representing environmentalists, commercial fishermen, recreational anglers, surfers, boaters, family farmers, faith communities, environmental justice advocates, labor unions, civic associations and recreational enthusiasts.

The copyrighted art on the cover was created by Sandra Noel of Vashon Island, Washington.

The text of this report is located on Clean Water Network's web page: <http://www.cwn.org>. Copies of this report are available for \$5.00 each, which includes tax and postage from:

Clean Water Network  
1200 New York Ave, Suite 400  
Washington D.C. 20005  
(Please make checks payable to NRDC)  
Phone 202/289-2395 x Fax 202/289-1060  
Email: [cleanwaternt@igc.org](mailto:cleanwaternt@igc.org)

[Table of Contents](#)

[Turn the page...](#)

# References

- Able, K.W., K.L. Heck, Jr., M.P. Fahay, and C.T. Roman. 1988. Use of salt-marsh peat reefs by small juvenile lobsters on Cape Cod, Massachusetts. *Estuaries* 11(2): 83-86.
- Alexander, C.E. and M.A. Broutman. 1996. An inventory of coastal wetlands of the United States. National Oceanic and Atmospheric Administration, Washington, D.C.
- Allen, L.G. 1988. Recruitment, distribution, and feeding habits of young-of-the-year California halibut (*Paralichthys californicus*) in the vicinity of Alamitos Bay-Long Beach Harbor, California, 1983-1985. *Bull. So. Calif. Acad. Sci.* 87(1), 1988, pp. 19-3
- Allen, L.G., R.E. Jensen and J.R. Sears. Open coast settlement and distribution of young-of-the-year California halibut (*Paralichthys californicus*) along the southern California coast between Point Conception and San Mateo Point, June-October, 1988. Pages 145-152 In C.W. Haugen (ed.) *The California halibut, Paralichthys californicus, resource and fisheries*. Calif. Dept. Fish and Game, Fish Bull. 174.
- Allen M.J. and K.T. Herbinson. 1990. Settlement of juvenile California halibut, *Paralichthys californicus*, along the coasts of Los Angeles, Orange, and San Diego Counties in 1989. *Calif. Coop. Ocean. Fish. Invest. Rep.* 31: 84-96.
- American Sportfishing Association. 1997. The 1996 Economic Impact of Sport Fishing in the United States. Prepared by Maharaj and Carpenter. December.
- Barrett, B.B., and M.C. Gillespie. 1973. Primary factors which influence commercial shrimp production in coastal Louisiana. *La. Dept. Wildl. Fish. Tech.* 9. 28 pp.
- Barshaw, D.E., and D.R. Bryant-Rich. 1988. A long-term study on the behavior and survival of early juvenile American lobster, *Homarus americanus* in three naturalistic substrates: eelgrass, mud and rocks. *Fish. Bull.* 86(4): 789-796.
- Boreman, J. 1983. Status of bluefish along the Atlantic coast, 1982. U.S. Nat. Mar. Fish. Serv., Woods Hole lab. Ref. Doc. No. 83-28. 36 pp.
- Bulger, A.J., B.P. Hayden, M.E. Monaco, D.M. Nelson, M.G. McCormick-Ray. 1993. Biologically-based estuarine salinity zones derived from a multivariate analysis. *Estuaries*, vol. 16, no. 2, pp. 311-322. Estuarine Research Federation.
- California Dept. of Fish and Game. Review of some California Fisheries for 1994. *CalCOFI Rep.*, Vol. 36, 1995. 12pp.
- California Legislature. 1988. Restoring the balance. Rpt. of the Adv. Cmte. on Salmon and Steelhead to the Joint Legislative Cmte. on Fisheries and Aquaculture, Sacramento. 84 pp.
- Carlson, H.R., R.E. Haight and K.J. Krieger. 1982. Species composition and relative abundance of demersal marine life in the waters of southeastern Alaska. 1969-1981. NWAFC Processed Rpt. 82-16. Nat. Mar. Fish. Serv., NOAA, Dept. of Commerce. 106 pp.
- Cato, J.C. and H.E. Kumpf. The economic influence of population growth, fisheries, coastal and marine industries, and tourism derived from use of the Gulf of Mexico. pp. 153-160, *Proc. of the Conference on the Environmental and Economic Status of the Gulf of Mexico*, Gulf of Mexico Program, New Orleans, LA.
- Cavit, N.H. 1981. Dependence of menhaden catch on wetlands habitats, a statistical analysis. U.S.

Fish Wildlife Service. National Geosystems Team. Unpublished report.

Chambers, J.R. 1991. Coastal degradation and fish population losses. In R.H. Stroud (ed.) Proc. of a Symposium on Coastal Fish Habitat Conservation, Baltimore, MD. Published by the National Coalition for Marine Conservation, Savannah, GA.

Christmas, J.Y., and R.S. Waller. 1975. Location and time of menhaden spawning in the Gulf of Mexico. Gulf Coast Research Laboratory. 20 pp.

Congressional Research Service. 1995. Wetlands Legislation: Comparison of Two Bills. Document 95-796 ENR.

Conrad, Earl. 1980. Gulf Stream North. Second Chance Press, Sag Harbor, N.Y.

Dobkin, S. 1961. Early developmental stages of pink shrimp, *Penaeus duorarum* from Florida waters. U.S. Fish Wildl. Serv. Fish Bull. 61: 321-349.

Douglas, Marjory Stoneman. 1947. The Everglades: River of Grass.

Dunbar, J.B., L.D. Britsch and E.B. Kemp III. 1992. Land loss rates, Louisiana coastal plain, report 3, technical report GL-90-2. U.S. Corps of Engineers.

Environmental Defense Fund and World Wildlife Fund. 1992. How wet is a wetland? The impacts of the proposed revisions to the federal wetlands delineation manual. 175 p. New York, N.Y. and Washington, D.C.

Fedler, A.J. and D.M. Nickum. The 1991 economic impact of sport fishing in the United States. Sports Fishing Institute, Washington, D.C., p. 2

Field, D.W., A.J. Reyer, P.V. Genovese and B.D. Shearer. 1991. Coastal wetlands of the United States, an accounting of a valuable national resource. Dept. of Commerce, NOAA, and Dept. of the Interior, U.S. Fish and Wildlife Service, Washington, D.C.

Fremling, C.R., J.L. Rasmussen, R.E. Sparks, S.P. Cobb, C.F. Bryan and T.O. Clafin. 1989. Mississippi River fisheries: a case history, pp. 309-351. In D.P. Dodge [ed.]. Proceedings on the International Large River Symposium. Can. Spec. Publ. Fish. Aquat. Sci. 106.

Friedland, K.D., G.C. Garman, A.J. Bejda, A.L. Studholme, and B. Olla. 1988. Interannual variation in diet and condition in juvenile bluefish during estuarine residency. Trans. Am. Fish. Soc. 117: 474-479.

Frey, H.W., ed. 1971. California's living marine resources and their utilization. Calif. Dept. Fish and Game. 148 pp.

Giles, J.H., and G. Zamora. 1973. Cover as a factor in habitat selection by juvenile brown (*Penaeus aztecus*) and white shrimp (*P. setiferus*). Trans. Am. Fish. Soc. 102(1): 144-145.

Guillory, V., J. Geaghan and J. Roussel. 1983. Influence of environmental factors on Gulf menhaden recruitment. Louisiana Dept. Wildlife and Fish., Tech. Bull. No. 37, 32 pp.

Gulf of Mexico Fishery Management Council. 1981. Fishery management plan for the shrimp fishery of the Gulf of Mexico, United States waters. Tampa, FL.

Gunter, G., and J.C. Edwards. 1969. The relation of rainfall and freshwater drainage to the production of the penaeid shrimps (*Penaeus fluviatilis* Say and *Penaeus aztecus* Ives) in Texas and Louisiana waters. FAO Fish. Rep. 57(3): 875-892.

Haaker, P.L. 1975. The biology of the California halibut, *Paralichthys californicus* (Ayers), in Anaheim Bay, California. pp. 137-151 In E.D. Lane and C.W. Hill, eds. The marine resources of Anaheim Bay. California Dept. Fish and Game, Fish Bull. 165.

Hersey, John. 1988. Blues. Random House, Inc. New York. 205 pp.

Heese, L.W., C.W. Wolfe, and N.K. Cole. 1988. Some aspects of energy flow in the Missouri River ecosystem and a rationale for recovery. In: North Central Division, American Fisheries Society, Spec. Pub. No. 8.

Heese, L.W. 1989. The changing Missouri River: a legacy for fish, wildlife and humankind. Pages 67-79 in J.A Kusler and M. Davidson, eds. Wetlands and river corridor management, proceedings of an international symposium, July 5-9, 1989, Charleston, South Carolina. Association for Wetland Managers, Bernes, N.Y.

Heese, L.W., J.C. Schmulbach, J.M. Carr, K.D. Keenlyne, D.G. Unkenholz, J.W. Robinson and G.E. Mestl. 1989. Missouri River fishery resources in relation to past, present, and future stresses, pp. 352-371. In D.P. Dodge [ed.]. Proceedings on the International Large River Symposium. Can. Spec. Publ. Fish. Aquat. Sci. 106.

Hofstra, Terrence D. 1983. Management alternatives for the Redwood Creek estuary. Redwood National Park, Arcata, Calif. 50 pp.

Hoss, Donald E. and Gordon W. Thayer. 1993. The importance of habitat to the early life history of estuarine dependent fishes. Pages 147-158, Proc. American Fisheries Society Symposium 14.

Hull, David. 1986. Restoring and protecting Humboldt Bay's aquatic resources. Pages 120-123 In Proc. of a Symposium on Managing Inflows to California's Bays and Estuaries, The Bay Institute, Sausalito, Calif.

June, F.C., and L.L. Chamberlin. 1959. The role of the estuary in the life history of the Atlantic menhaden. Proc. Gulf and Carib. Fish. Inst., pp.41-45.

Louisiana Department of Wildlife and Fisheries. 1997. The Economic Benefits of Fisheries, Wildlife and Boating Resources in the State of Louisiana. March. Southwick Associates.

Kier, William M. 1992. The relationship between streamflow, opossum shrimp habitat and coho salmon growth and survival in the Lagunitas Creek estuary. Expert testimony presented to the California State Water Resources Control Board, hearing of May 12, 1992 concerning water use in Lagunitas Creek, Marin County, CA.

Knapp, G. and T. Smith. 1991. Seafood industry sector report to the Alaska Dept. of Commerce and Economic Development. Institute of Social and Economic Research, Univ. of Alaska, Anchorage. 162 pp.

Kramer, S.H. 1990. Distribution and abundance of juvenile California halibut, *Paralichthys californicus*, in shallow waters of San Diego County. Pages 99-126 In C.W. Haugen (ed.) The California halibut, *Paralichthys californicus*, resources and fisheries. Calif. Dept. Fish and Game, Fish Bull. 174.

Kramer, S.H. and J.R. Hunter. 1987. Southern California wetland and shallow water habitat investigation, annual report for the fiscal year 1987. U.S. Dept. Commerce, NOAA, NMFS, SWFC, La Jolla, Calif., 12 pp.

Kutkuhn, J.H. 1966. The role of estuaries in the development and perpetuation of commercial shrimp resources. Pages 16-36 In R.F. Smith, A.H. Swartz, and W.H. Massman, eds. A symposium on estuarine fisheries. Am. Fish. Soc. Spec. Publ. 3.

McKeon, Joseph F. 1985. Downstream migration, growth, and condition of juvenile fall chinook salmon in Redwood Creek, Humboldt County, California. M.S. thesis, Humboldt State University, Arcata, Calif. 90 pp.

Merrell, Theodore R. and K.V. Koski. Habitat values of coastal wetlands for Pacific coast salmonids.

Pages 256-266 In P.H. Greeson et al. (eds.) Wetland functions and values: the state of our understanding. Proc. National Symposium on Wetlands, American Water Resources Association, Minneapolis, MN.

Monaco, M.E., T.A. Lowery and R.L. Emmett. 1992. Assemblages of U.S. west coast estuaries based on the distribution of fishes. *Jour. of Biogeography*, vol. 19, pp. 251-267.

Muncy, R.J. 1984. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Gulf of Mexico) — white shrimp. U.S. Fish and Wildlife Service FWS/OBS-82/11.20 U.S. Army Corps of Engineers, TR EL-82-4. 19 pp.

National Research Council. 1992. Restoration of Aquatic Ecosystems. National Academy Press. Washington, D.C. 552 pp.

Norcross, J.J., S.L. Richardson, W.H. Massman, and E.B. Joseph. 1974. Development of young bluefish (*Pomatomus saltatrix*) and distribution of eggs and young in Virginia coastal waters. *Trans. Am. Fish. Soc.* 103: 477-497.

Odum, W.E. 1971. Pathways of energy flow in a south Florida estuary. Ph.D. Dissertation. University of Miami, Fla. Diss. Abstr. Univ. Mich. 70-18, 156: 1898B.

Perret, W.S. 1968. Menhaden or pogies: Louisiana's most valuable commercial fish. *Louisiana Conserv.* 20(1 and 2): 14-15.

Perret, W.S. and M.F. Chatry. 1991. Coastal Louisiana - abundant renewable resources in peril. In: *Coastal wetlands*, S.H. Bolton (ed.), pp. 317-331. American Society of Civil Engineers Press, New York, N.Y.

Perret, W.S. and W.E. Schaaf. 1991. Empirical model of the trophic basis for fishery yield in coastal waters of the eastern USA. *Trans. American Fish. Soc.*, 120:459-473, 1991.

———. 1992. Atlantic marsh-estuarine nearshore detrital system (amends) model. In K.W. Turgeon (ed.) *Proc. of a workshop on marine ecosystem modeling*. Dept. of Commerce, NOAA. Frederick, MD. April 6-8, 1992.

Perret, W.S., J.E. Roussel, J.F. Burdon and J.F. Pollard. 1993. Long term trends of some trawl-caught estuarine species in Louisiana. In: *Proc. of the 8th symposium on coastal and ocean management*, O.T. Magoon, W.S. Wilson, H. Converse and L.T. Tobin eds.

Peters, David S. and Ford A. Cross. 1992. What is coastal fish habitat? Pages 17-22 In Richard H. Stroud (ed.), *Stemming the tide of coastal fish habitat loss*. Proc. of a Symposium on Conservation of Coastal Fish Habitat, Baltimore, MD. Published by the National Coalition for Marine Conservation, Inc., Savannah, GA.

Plummer, K.E., E.E. DeMartini, and D.A. Roberts. 1983. The feeding habits and distribution of juvenile-small adult California halibut (*Paralichthys californicus*) in coastal waters off northern San Diego County, *CalCOFI Rep.*, 24: 194-201.

Powell, A.B., D.E. Hoss, W.F. Hettler, D.S. Peters, and S. Wagner. 1989. Abundance and distribution of ichthyoplankton in Florida Bay and adjacent waters. *Bull. of Marine Sci.*, 44(1), p. 35-48. 1989.

Rasmussen, J.L. 1997. American Fisheries Society Position Statement – Floodplain Management. [Http:// www.esd.ornl.gov/societies/AFS/ps-flood.html](http://www.esd.ornl.gov/societies/AFS/ps-flood.html)

Reid, G.K. 1957. Biologic and hydrologic adjustment in a disturbed gulf coast estuary. *Limnol. and Oceanogr.* 2: 198-212.

Reimers, P.E. 1973. The length of residence of juvenile fall chinook salmon in Sixes River, Oregon. *Research Reports of the Fish Commission of Oregon.* (4)2.

Robison, Clayton D., Jr. 1993. Alaska wetlands are not different. National Wetlands Newsletter, Environ. Law. Instit., Vol. 15, No. 5.

Rozenfurt, M.A. and I. Haydock. 1992. Effects of fresh water development and water pollution policies on the world's river, delta, estuary, coastal zone ecosystems. In H. S. Bolton, ed. Coastlines of the World, Amer. Soc. of Civil Eng., New York.

Simenstad, C.A., K.L. Fresh and E.O. Salo. 1982. The role of Puget Sound and Washington coastal estuaries in the life history of Pacific salmon: an unappreciated function. Pages 343-364, In Victor S. Kennedy ed. Estuarine Comparisons, Academic Press, New York, New York. 704 pp.

Skinner, John E. 1962. An historical review of the fish and wildlife resources of the San Francisco Bay area. Wat. Proj. Branch Rpt. 1, Calif Dept of Fish and Game, Sacramento. 225 pp.

Smith, Jerry J. 1990. The effects of sandbar formation and inflows on aquatic habitat and fish utilization on Pescadero, San Gregorio, Waddell and Pomponio Creek estuary/lagoon systems, 1985-1989. San Jose State Univ., Dept. of Biol. Sci. Interagency Agreement Rpt. 84-04-324.

Spratt, Jerome D. 1981. Status of the Pacific herring, *Clupea harengus pallasii*, resource in California 1972 to 1980. Fish Bull. 171, Calif. Dept. of Fish and Game, Sacramento. 107 pp.

Trent, L., E.J. Pullen, and R. Proctor. 1976. Abundance of macro-crustaceans in a natural marsh and marsh altered by dredging, bulkheading, and filling. U.S. Natl. Mar. Fish. Serv. Fish Bull. 74(1): 195-200.

Turner, R.E. 1977. Intertidal vegetation and commercial yields of penaeid shrimp. Trans. Am. Fish. Soc. 106(5): 411-416.

U.S. Dept. of Commerce. 1992. Our living oceans, report on the status of U.S. living marine resources, 1992. NOAA Tech. Memo. NMFS-F/SPO-2. Washington, D.C. 148 pp.

———. 1994. National Oceanic and Atmospheric Administration, Estuarine living marine resources program (ELMR), draft species life history notes for the North Atlantic Region. NOS/ORCA/SEA Biogeographic Characterization Branch, Silver Spring, MD.

———. 1996. Fisheries of the United States, current fisheries statistics no. 9600. Washington, D.C.

U.S. Dept. of Interior. USFWS News Release, April 30, 1997. Billions and Billions Spent: Wildlife Related Recreation Continues to be National Economic Force.

———. USFWS News Release, August 27, 1997. Service Releases Preliminary State-by-State Report on Wildlife Related Recreation.

———. Draft Environmental Impact Statement for Proposed Expansion of Big Muddy National Fish and Wildlife Refuge, Missouri. 1997. USFWS Puxico, MO. 102 p. and appendices.

U.S. Dept. of Interior and U.S. Dept. of Commerce. 1993. 1991 national survey of fishing, hunting and wildlife-associated recreation. U.S. Govt. Printing Office, Washington, D.C.

U. S. Department of the Interior, Fish and Wildlife Service and U. S. Department of Commerce, Bureau of the Census. 1996 National Survey of Fishing, Hunting and Wildlife-Associated Recreation. Washington, D. C. 1997.

U.S. Environmental Protection Agency (EPA). 1992. Status and trends report on aquatic resources in the San Francisco Estuary. San Francisco Estuary Project. Oakland, CA. 257 pp. plus app's.

———. 1998. Chesapeake Bay Program Phase IV Watershed Model, 1985 Reference Scenario, per Chesapeake Bay Commission.

\_\_\_\_\_. 1998. Clean Water Action Plan: Restoring and Protecting America's Waters. Washington, D.C. Doc. No. EPA-840-R-98-001.

\_\_\_\_\_. 1990. Progress report for the baywide nutrient reduction strategy. Region III, Chesapeake Bay Program Office, Annapolis, Md.

U.S. House of Representatives. 1997. H.R. 2762, October 29, 1997. A Bill to amend the Federal Water Pollution Control Act to improve the protection of the Nation's wetlands and watersheds, and for other purposes. 105th Congress, 1st Session.

Viosca, P. 1928. Louisiana wetlands and the value of their wildlife and fishery resources. *Ecol.* 9(2):216-229.

Wild, Paul W. and Robert N. Tasto, eds. 1983. Life history, environment, and mariculture studies of the Dungeness crab, *Cancer magister*, with emphasis on the central California fishery resource. *Fish Bull.* 172, Calif. Dept. of Fish and Game, Sacramento. 352 pp.

Wilk, S.J. 1977. Biological and fisheries data on bluefish, *Pomatomus saltatrix*. U.S. Nat. Mar. Fish. Serv., Sandy Hook Lab., Highlands, N.J. Tech. Serv. Rep. 11.

Williams, A.B. 1958. Substrates as a factor in shrimp distribution. *Limnol. Oceanogr.* 3(3): 283-290.

## [Table of Contents](#)