


Recovery Plan for the Kootenai River Population of the White Sturgeon

(Acipenser transmontanus)



**Recovery Plan for the White Sturgeon (*Acipenser
transmontanus*): Kootenai River Population**

Published by
Region 1
U.S. Fish and Wildlife Service
Portland, Oregon

Approved: Anne Badgley
Regional Director, Region 1, U.S. Fish and Wildlife Service

Date: 9/30/99

DISCLAIMER PAGE

Recovery plans delineate reasonable actions that are believed necessary to recover or protect the species. Plans are prepared by the U.S. Fish and Wildlife Service, sometimes with the assistance of recovery teams, contractors, State agencies, Tribal agencies, and others. Objectives will be attained and any necessary funds made available subject to budgetary and other constraints affecting the parties involved, as well as the need to address other priorities. Recovery plans do not necessarily represent the views nor the official positions or approval of any individuals or agencies involved in the plan formulation, other than the U.S. Fish and Wildlife Service. Recovery plans represent the official position of the U.S. Fish and Wildlife Service *only* after they have been signed by the Director or Regional Director as *approved*. Approved recovery plans are subject to modification as dictated by new findings, changes in species status, and the completion of recovery tasks.

Literature Citation: U.S. Fish and Wildlife Service. 1999. Recovery Plan for the White Sturgeon (*Acipenser transmontanus*): Kootenai River Population. U.S. Fish and Wildlife Service, Portland, Oregon. 96 pp. plus appendices.

Additional copies may be purchased from:

Fish and Wildlife Reference Service
5430 Grosvenor Lane, Suite 110
Bethesda, Maryland 20814-2158
301-492-6403 or 1-800-582-3421
E-mail: fwrs@mail.fws.gov
<http://fa.r9.fws.gov/r9fwrs/>

The fee for the Plan varies depending on the number of pages of the Plan.

ACKNOWLEDGMENTS

Cover art for this plan was graciously provided by Calvin R. Sprague. The Recovery Plan was prepared by the Kootenai River White Sturgeon Recovery Team comprising the following individuals: ¹

Kootenai River White Sturgeon Recovery Team

Steve Duke

Team Leader
U.S. Fish and Wildlife Service
Snake River Basin Office
1387 S. Vinnell Way, Suite 368
Boise, Idaho 83709

Paul Anders

University of Idaho
Fish Genetics Lab
Aquaculture Research Institute
Moscow, Idaho 83844

Gordon Ennis

Canada Dept of Fisheries and Oceans
555 West Hastings Street
Vancouver, British Columbia
Canada V6B 5G3

Robert Hallock, PhD

U.S. Fish and Wildlife Service
Upper Columbia Basin Office
11103 East Montgomery Dr., #2
Spokane, Washington 99206

Jay Hammond

British Columbia Ministry of
Environment, Lands and Parks
401-333 Victoria Street
Nelson, British Columbia
Canada V1L 4K3

Sue Ireland

Kootenai Tribe of Idaho
P.O. Box 1269
Bonners Ferry, Idaho 83805

Jeff Laufle

U.S. Army Corps of Engineers
P.O. Box 3755
Seattle, Washington 98124

Larry Lockard

U.S. Fish and Wildlife Service
Ecological Services
780 Creston Hatchery Road
Kalispell, Montana 59901

Brian Marotz

Montana Fish, Wildlife & Parks
490 North Meridian Road
Kalispell, Montana 59901

Vaughn L. Paragamian

Idaho Department of Fish and Game
2750 W. Kathleen Ave.
Coeur d'Alene, Idaho 83814

Scott Bettin

Bonneville Power Administration
P.O. Box 3621-EWI
Portland, Oregon 97208

¹ Ray Lauzier, Canada DFO and Rick Westerhof, BPA, were members of the recovery team during development of the draft recovery plan.

EXECUTIVE SUMMARY

Recovery Plan for the White Sturgeon (*Acipenser transmontanus*): Kootenai River Population

Current Species Status: The Kootenai River population of white sturgeon was listed as endangered on September 6, 1994 (59 FR 45989). This white sturgeon population has been in general decline since the mid-1960's. In 1997 the population was estimated to be approximately 1,468 wild fish with few individuals less than 25 years of age. In 1997, the wild population was augmented with the release of 2,283 juvenile white sturgeon reared in the Kootenai Tribal hatchery in Bonners Ferry, Idaho.

Habitat Requirements and Limiting Factors: The Kootenai River population of white sturgeon became isolated from other white sturgeon in the Columbia River basin during the last glacial age (approximately 10,000 years ago). Once isolated, the population adapted to the predevelopment habitat conditions in the Kootenai River drainage. Historically, spring runoff peaked during the first half of June in the Kootenai River upstream of the existing Libby Dam in Montana. Runoff from lower elevations between Libby Dam and Bonners Ferry, Idaho was somewhat earlier, peaking in late May. Combined flows were often in excess of 1,700 cubic meters per second (m^3/s) (60,000 cubic feet per second (cfs)). During the remainder of the year, river flows declined to basal conditions of 113 to 226 cubic meters per second (4,000 to 8,000 cubic feet per second). Annual flushing events re-sorted river sediments providing a clean cobble substrate conducive to insect production and sturgeon egg incubation. Side channels and low-lying deltaic marsh lands were undiked at this time, providing productive, low velocity backwater areas. Nutrient delivery in the system was unimpeded by dams and occurred primarily during spring runoff. Flood plain ecosystems like the predevelopment Kootenai River are characterized by seasonal floods that promote the exchange of nutrients and organisms in a mosaic of habitats and thus enhance biological productivity (Bayley 1995; Junk et al. 1989; Sparks 1995).

Modification of the Kootenai River white sturgeon's habitat by human activities has changed the natural hydrograph of the Kootenai River, altering white sturgeon spawning, egg incubation, and rearing habitats; and reducing overall biological productivity. These factors have contributed to a general lack of recruitment in the white sturgeon population since the mid-1960's.

Recovery Objectives: Downlisting and Delisting. The short-term recovery objectives are to re-establish successful natural recruitment and prevent extinction through the use of conservation aquaculture. The long-term objective is to downlist and then delist the fish when the population becomes self-sustaining.

Recovery Criteria: Criteria required for reclassification or downlisting to threatened status include:

1. Natural production of white sturgeon occurs in at least 3 different years of a 10-year period; a naturally produced year class is demonstrated when at least 20 juveniles from a year class are sampled at more than 1 year of age.
2. The estimated white sturgeon population is stable or increasing and juveniles reared through a conservation aquaculture program are available to be added to the wild population each year for a 10-year period. Each of these year classes must be large enough to produce 24 to 120 sturgeon surviving to sexual maturity.
3. A long-term Kootenai River Flow Strategy is developed in coordination with interested State, Federal, and Canadian agencies and the Kootenai Tribe at the end of the 10-year period based on results of ongoing conservation efforts, sturgeon habitat research, and fish productivity studies. An important element of this strategy is demonstration of the repeatability of in-stream environmental conditions necessary to produce recruits (as described above) in future years.

Specific delisting recovery criteria have not been identified at this time, but will be developed as new population status, life history, biological productivity, and flow augmentation monitoring information is collected. However, recovery will not be complete until there is survival to maturity and natural reproduction of juvenile white sturgeon added to the wild population from the conservation aquaculture program. This may take upwards of 25 years since that is the approximate period for juvenile female white sturgeon to reach sexual maturity and reproduce to complete a new generation or spawning cycle.

Actions Needed:

- o Identify and restore white sturgeon habitats necessary to sustain white sturgeon reproduction (spawning and early age recruitment) and rearing while minimizing impacts on other uses of Kootenai River basin waters.

- o Develop and implement a conservation aquaculture program to prevent the extinction of Kootenai River white sturgeon. The conservation aquaculture program will include protocols on broodstock collection, propagation, juvenile rearing, fish health, genetics, and stocking.
- o Work within operational guidelines for Libby Dam based upon Kootenai Integrated Rule Curves (KIRC) developed by Montana Fish, Wildlife, and Parks to balance white sturgeon recovery with requirements for other aquatic species and recreational fisheries within the Kootenai River drainage, and VARQ (an enhanced flood control protocol), to ensure that more water is available for white sturgeon, salmon, and all species in lower water years.
- o Continue research and monitoring programs (with achievable and measurable objectives) on life history, habitat requirements for all life stages, population status, and trends of the Kootenai River white sturgeon.
- o Protect Kootenai River white sturgeon and their habitats using available regulatory mechanisms.
- o Evaluate how changes in biological productivity in the Kootenai River basin affect white sturgeon and their habitats.
- o Evaluate the effects of contaminants and possible additional biological threats, e.g. predation and species composition, on Kootenai River white sturgeon and their habitats.
- o Increase public awareness of the need to protect and recover Kootenai River white sturgeon.
- o Balance white sturgeon recovery measures with requirements for other aquatic species and recreational fisheries within the Kootenai River drainage.
- o Secure funding for implementation of recovery tasks.

Estimated Cost of Recovery : Costs for some tasks are estimated to be \$7,456,000 for the first 5 fiscal years. Total estimated recovery costs will likely increase as new information is received and as the ongoing biological studies are completed. Estimated costs do not include costs associated with native fish monitoring tasks. Future total costs may also decrease as some research tasks are completed.

Other Physical and Economic Impacts from Recovery: Implementing many of the conservation actions proposed in this recovery plan will create additional economic or environmental impacts, as well as associated benefits, not normally considered in estimating the “costs” of recovery. Economic or environmental impacts may include foregone power generation opportunities, reduced flood control, and possibly negative impacts to other regional resident fish.

Associated benefits include the partial restoration of a more natural Kootenai River hydrograph and flood plain function that benefits resident fish and wildlife. Periodic flushing flows would cleanse Kootenai River gravels and improve aquatic insect production. Improving aquatic ecosystem health leading to improved regional fisheries will provide secondary economic benefits to local communities. Such benefits go beyond the “benefits” typically considered in recovery actions. Conversely, failure to implement proposed recovery actions would have hidden costs that are typically not considered in cost/benefit analysis.

Date of Recovery: At a minimum, at least 25 years following implementation of an approved recovery plan are necessary before delisting of the white sturgeon population can be considered. This 25-year period would allow juveniles added to the population in the first 10 years to reach maturity and begin reproducing a new generation.

What is a recovery plan? A recovery plan is a template for the recovery of threatened or endangered species and their habitats. The recovery plan describes the process by which the decline of a listed species may be reversed and known threats to its long-term survival can be removed. Therefore, recovery is the restoration of a listed species to the point where they become secure, self-sustaining components of their ecosystem.

An approved recovery plan is not a decision document but is intended to provide information and guidance that the U.S. Fish and Wildlife Service believes will lead to recovery of a listed species, including its habitat. The recovery plan provides information necessary to describe the current status of the listed species as well as on-going or proposed actions designed to aid in the species ultimate recovery. Many of the recovery actions (or tasks) in this document will require further environmental analysis and public review, especially those actions taken by Federal agencies.

This final recovery plan serves as a guidance document listing various conservation actions for the recovery of the white sturgeon population within the Kootenai River basin and the ecosystem upon which it depends. It was developed by a recovery team composed of persons from State, Federal, Tribal, and Canadian agencies who have experience with this population of white sturgeon or the threats it faces. Because the white sturgeon population is only one component of its ecosystem, the recovery team took a holistic approach that will address other sensitive aquatic species that are dependent upon the Kootenai River drainage. Efforts proposed for Kootenai River white sturgeon recovery should benefit many other native aquatic species and possibly aid the restoration of declining species in Kootenai River drainage habitats before their status becomes critical. However, actions that will directly benefit the white sturgeon are given highest priority. Other lower priority actions, which could benefit nonlisted aquatic species and further contribute to overall ecosystem recovery, are also included in the recovery plan.

What is the Kootenai River ecosystem? An ecosystem is defined as an ecological community that together with its environment, functions as a unit. For the purposes of this recovery plan, the Kootenai River ecosystem is defined as the habitat and aquatic species complex within the Kootenai drainage basin including Koocanusa Reservoir upstream of Libby Dam, Kootenai River downstream including tributary streams, backwater sloughs, deltaic marshlands, and Kootenay Lake in British Columbia downstream to Corra Linn Dam at the outlet of the West Arm of Kootenay Lake. (Kootenai is spelled Kootenay in Canada.)

TABLE OF CONTENTS

	<u>Page</u>
EXECUTIVE SUMMARY	iii
WHAT IS A RECOVERY PLAN?	vii
WHAT IS THE KOOTENAI RIVER ECOSYSTEM?	vii
GLOSSARY	xii
LIST OF SYMBOLS AND ABBREVIATIONS	xvii
PART I. INTRODUCTION	
A. Overview	1
B. General Description	1
C. Aquatic Community	6
D. Reasons for Decline	7
E. Conservation Measures	11
F. Strategy for Recovery	34
PART II. RECOVERY	
A. Recovery Objectives	37
B. Recovery Criteria	37
C. Recovery Measures Narrative	40
D. References Cited	84

	<u>Page</u>
PART III. IMPLEMENTATION SCHEDULE	88
PART IV. APPENDICES	96
Appendix A. Current fish fauna of the Kootenai River basin.	
Appendix B. Examination of the effects of two alternative flow augmentation strategies on the Kootenai River ecosystem.	
Appendix C. Kootenai basin integrated rule curves and tiered approach for white sturgeon flow release from Libby reservoir.	
Appendix D. Breeding plan to preserve the genetic variability of the Kootenai River white sturgeon.	
Appendix E. White sturgeon broodstock collection protocols.	
Appendix F. Summary of the agency and public comments on the draft recovery plan for Kootenai River white sturgeon.	

LIST OF FIGURES

	<u>Page</u>
1. Map of the Kootenai River basin.	2
2. Map of the Kootenai River basin in Idaho and Montana.	5
3. Mean monthly Kootenai River flows at Bonners Ferry for 1928-1972 (pre-Libby Dam) and 1973-1995 (post-Libby Dam) periods.	9
4. Kootenai River estimated flows and water temperature observed during April through July, 1991.	12
5. Kootenai River estimated flows and water temperature observed during April through July, 1992.	14
6. Kootenai River estimated flows and water temperature observed during April through July, 1993.	15
7. Kootenai River estimated flows and water temperature observed during April through July, 1994.	17
8. Kootenai River estimated flows and water temperature observed during April through July, 1995.	20
9. Kootenai River estimated flows and water temperature observed during April through July, 1996.	22
10. Kootenai River estimated flows and water temperature observed during April through July, 1997.	23
11. Flow chart summarizing proposed Kootenai River white sturgeon recovery measures.	41

LIST OF TABLES

	<u>Page</u>
1. Summary of historical harvest regulations for white sturgeon within the Kootenai River drainage in the United States and Canada.	31
2. “Tiered” volumes of water for sturgeon flow enhancement to be provided to Bonners Ferry according to the April-August volume runoff forecast at Libby.	48

GLOSSARY

Anecdotal Evidence	Information passed along by word of mouth but not documented scientifically.
Bedload	Streambed materials that are washed downstream and redeposited in a new location.
Biological Productivity	A measure of growth in living systems.
Biological Trophic Levels	Steps in the food chain from plants through plant eaters to meat eaters.
Biomass	The total weight of a living organism or a population of organisms.
Chlorinated Biphenyls	A contaminant that accumulates in the fatty tissues of organisms that can cause health problems.
Community Respiration	The amount of energy used by all of the organisms in a specified locality.
Conservation Aquaculture	A hatchery-based, captive culture program designed to 1) preserve the Kootenai River white sturgeon gene pool (genetic variation) and 2) rebuild the natural age class structure of white sturgeon in the wild through the release of hatchery-reared juvenile fish. The program is based on a breeding plan that includes protocols on adult broodstock collection, hatchery spawning and rearing, fish health, and genetics.

Delta (as in tributary)	Streambed materials that accumulate near the mouth of a stream.
Discharge	Water flow volume, usually used to describe a volume released from a dam.
Electrophoretic Analysis	A laboratory technique to examine genetic differences between similar species. Protein samples are placed in an electrical field producing bands on a gel plate. The bands are used like fingerprints to distinguish genetic traits.
Empirical Data	Information derived from measurements made in "real life" situations (e.g. field data).
Flow Ramping	The act of creating a gradual rather than abrupt change in flow. Typically used to define allowable fluctuations below a hydropower dam.
Gas supersaturation	Aquatic conditions that result from turbulence that allows water to absorb nitrogen or oxygen from air bubbles trapped several feet below the surface. As these waters rise back to the surface, they become supersaturated because pressure drops. Some of these gases may become trapped in a fish's blood vessels and cause injury or death.
Habitat Use Curve	A graph describing the distribution/occurrence of fish over a range of a specific environmental variable (e.g. velocity, temperature or depth).

Hydrograph	The recorded variations in stream discharge over time. Useful when comparing effects and changes in stream flow and depth between average natural conditions and altered stream flows (i.e. from dams and diversions).
In-stream Flow Incremental Methodology (IFIM)	A process that uses river channel measurements and hydraulic characteristics to estimate the amount of available fish habitat under various river discharges.
Koocanusa Reservoir	Also known as Libby Reservoir or Lake Koocanusa, located upstream of Libby Dam.
Kootenay Lake	A natural lake in British Columbia, which is regulated by Corra Linn Dam. The Kootenai River, downstream of Libby Dam enters Kootenay Lake from the south.
Limnological (limnology)	The science of the properties of fresh water including water chemistry, density, stratification and physical effects on living organisms.
Load Following	Short-term changes in hydropower operations to respond to subtle shifts in power demand. Flow fluctuations caused by load following are usually less dramatic than power peaking.
Microhabitat	Detailed description of where an animal lives.
Nutrient Dynamics	The way nutrients are used and reused, over time and distance, in a biological system.

Organochlorides	Complex toxic molecule containing carbon and chlorine that is soluble in fatty tissues and can cause health problems.
Photoperiod	A measurement of time exposed to light in a given day or series of days.
Power Peaking	Hydropower operations that occur for short time periods. Typically more power is generated during the day than at night, causing changes in stream flows.
Redox Potential	A measurable electric charge (volts) created when an oxidizing agent pulls electrons away from a reducing agent. This action is an important factor in nutrient cycling in water.
Recruitment	Survival of juveniles until they become a member of the spawning population.
Relative Abundance	A comparison of the number in one category to another (e.g. number of one species to another, male to female, young to old, etc.). Typically expressed as a percentage or proportion.
Reservoir Drawdown	Removing water from a reservoir and lowering the surface elevation.
Scutes	Hard ridges or bony structures along the back of sturgeons.
Tributary	A small stream or river, which enters and increases the volume of the receiving river, lake, or reservoir.
Vermiculite	A mineral mined from the earth having fire retardant and insulating properties.

Year class

All individuals of a fish population
spawned and hatched in a given year.

LIST OF SYMBOLS AND ABBREVIATIONS

Act	Endangered Species Act of 1973 (as amended)
B.C.	British Columbia
B.A.	Biological Assessment
B.O.	Biological Opinion
BPA	Bonneville Power Administration
BR	Bureau of Reclamation
cfs	cubic feet per second
C.I.	confidence interval
cm	centimeter
DFO	Canada Department of Fisheries and Oceans
FCRPS	Federal Columbia River Power System
FWS	U.S. Fish and Wildlife Service
IDFG	Idaho Department of Fish and Game
in	inch
IRC	Integrated Rule Curves
kg	kilogram
KIRC	Kootenai Integrated Rule Curves
km	kilometer
KRBSC	Kootenai River Basin Steering Committee
KTOI	Kootenai Tribe of Idaho
lb	pound
MELP	British Columbia Ministry of Environment, Lands and Parks
mi	mile
MFWP	Montana Department of Fish, Wildlife, and Parks
m ³ /s	cubic meters per second
NMFS	National Marine Fisheries Service
NPPC	Northwest Power Planning Council
PCB	polychlorinated biphenyl
ppm	parts per million
Program	Columbia River Basin Fish and Wildlife Program
PSMFC	Pacific States Marine Fisheries Commission
rkm	river-kilometer

rm
Service
U.S.
USACE

river-mile
U.S. Fish and Wildlife Service
United States
U.S. Army Corps of Engineers

PART 1 - INTRODUCTION

A. Overview

On September 6, 1994, the U.S. Fish and Wildlife Service listed the Kootenai River population of white sturgeon as an endangered species (59 FR 45989) under the authority of the Endangered Species Act of 1973, as amended.

The Kootenai River population is one of several land-locked populations of white sturgeon found in the Pacific Northwest. Although officially termed and listed as the “Kootenai River population of white sturgeon”, this white sturgeon population inhabits and migrates freely in the Kootenai River from Kootenai Falls in Montana downstream into Kootenay Lake, British Columbia, Canada (Figure 1).

The Endangered Species Act specifies that recovery plans should, to the maximum extent practicable, give priority to those listed species most likely to benefit from recovery actions. The recovery priority for the Kootenai River population of white sturgeon is 3C indicating that: 1) taxonomically, it is a "distinct population segment" of a species; 2) it is subject to a high degree of threat; 3) the recovery potential is high; and 4) the degree of potential for conflict with construction or other development projects is high.

B. General Description

White sturgeon (*Acipenser transmontanus*) occur along the Pacific coast from the Aleutian Islands to central California. In unimpounded river systems, the species migrates between the sea and fresh water, and reproduces in at least three large river systems: the Sacramento-San Joaquin River in California, the Columbia River basin in the Pacific Northwest, and the Fraser River system in British Columbia. The Kootenai River population of white sturgeon is one of 18 land-locked populations of white sturgeon found in the Pacific Northwest. Their distribution extends from Kootenai Falls, Montana, located 50 river-kilometers [rkm] (31 river-miles [rm]) below Libby Dam, downstream through Kootenay Lake to Corra Linn Dam on the lower West Arm of Kootenay Lake, British Columbia.




Figure 1. Map of the Kootenai River Basin.

Kootenai Falls may represent an impassible natural barrier to the upstream migration of white sturgeon although anecdotal evidence suggests the historic presence of white sturgeon upstream from Kootenai Falls in Montana and British Columbia. A natural barrier at Bonnington Falls downstream of Kootenay Lake has isolated the Kootenai River white sturgeon from other white sturgeon populations in the Columbia River basin since the last glacial age, approximately 10,000 years ago (Northcote 1973).

White sturgeon are included in the family Acipenseridae, which consists of 4 genera and 24 species of sturgeon. Eight species of sturgeon occur in North America with white sturgeon being one of five species in the genus *Acipenser*. White sturgeon were first described by Richardson in 1863 from a single specimen collected in the Columbia River near Fort Vancouver, Washington (Scott and Crossman 1973). White sturgeon are distinguished from other *Acipenser* by the specific arrangement and number of scutes (bony plates) along the body (Scott and Crossman 1973). The largest white sturgeon on record, weighing approximately 682 kilograms (1,500 pounds), was taken from the Snake River near Weiser, Idaho in 1898 (Simpson and Wallace 1982). Scott and Crossman (1973) describe a white sturgeon reported to weigh over 818 kilograms (1,800 pounds) from the Fraser River near Vancouver, British Columbia, date unknown. Individuals in landlocked populations tend to be smaller. The largest white sturgeon reported from the Kootenai River basin is a 159 kilograms (350 pounds) individual estimated at 85 to 90 years of age captured in Kootenay Lake during September 1995 (Lindsay 1995). White sturgeon are generally long-lived, with females living from 34 to 70 years (PSMFC 1992).

The size or age at first maturity for white sturgeon in the wild is quite variable (PSMFC 1992). In the Kootenai River system, females have been documented to mature as early as age 22 and males at age 16 (Paragamian et al. 1997). Only a portion of adult white sturgeon are reproductive or spawn each year, with the spawning frequency for females estimated at 2 to 11 years (PSMFC 1992). Spawning occurs when the physical environment permits egg development and cues ovulation. White sturgeon are broadcast spawners, releasing their eggs and sperm in fast water. Based upon recent studies, Kootenai River white sturgeon spawn during the period of historical peak flows from May through July

(Apperson and Anders 1991; Marcuson 1994). Spawning at peak flows with high water velocities disperses and prevents clumping of the adhesive eggs. Following fertilization, eggs adhere to the river substrate and hatch after a relatively brief incubation period of 8 to 15 days, depending on water temperature (Brannon et al. 1984). Recently hatched yolk-sac larvae swim or drift in the current for a period of several hours and then settle back into interstitial spaces in the substrate. Larval white sturgeon require an additional 20 to 30 days to metamorphose into juveniles with a full complement of fin rays and scutes.

Historically (pre-Libby Dam construction and operation), spawning areas for white sturgeon were not specifically known. White sturgeon monitoring programs conducted from 1990 through 1995 revealed that white sturgeon spawned within a 19 river-kilometer (12 river-mile) stretch of the Kootenai River, primarily from Bonners Ferry downstream to the lower end of Shorty's Island (Figure 2).

White sturgeon in the Kootenai River system and elsewhere are considered opportunistic feeders. Partridge (1983) found white sturgeon more than 70 centimeters (28 inches) in length feeding on a variety of prey items including clams, snails, aquatic insects, and fish. Andrusak (MELP, pers. comm., 1993) noted that kokanee (*Oncorhynchus nerka*) in Kootenay Lake, prior to a dramatic population crash beginning in the mid-1970's, were once considered an important prey item for adult white sturgeon.

Partridge (1983) noted that white sturgeon recruitment was intermittent and possibly decreasing from the mid-1960's to 1974 when Libby Dam started operations. This is demonstrated by the absence of white sturgeon year classes in samples collected in the early 1980's (i.e. 1965 to 1969, 1971, and 1975). Partridge speculated that the lack of recruitment in certain years was due in part to (1) the elimination of rearing areas for juveniles through diking of slough and marsh side-channel habitats; and (2) the increase in chemical pollutants, e.g. copper and zinc, released in the past from mineral processing facilities, which may have affected spawning or recruitment success.

Previous estimates of population size suggested that the Kootenai River white sturgeon population had declined from an estimated 1,194 fish in 1982 (Partridge




Figure 2. Map of the Kootenai River Basin in Idaho and Montana. Notable geographic features include Kootenai Falls, the suspected upstream migration barrier for white sturgeon, and the Kootenai River reach from Bonners Ferry downstream to Shorty's Island where white sturgeon spawning has been detected in recent years.

1983) to approximately 880 fish by 1990 (Apperson and Anders 1991). More recently, a refined white sturgeon population analysis using capture information collected from the Kootenai River and Kootenay Lake over a 4-year period estimated 1,468 adult fish (95 percent confidence interval: 740 to 2,197) and 87 wild juveniles. Although this revised estimated population is higher than the level when the white sturgeon was listed in 1994, the unbalanced population structure and primary factors affecting the listing decision persist.

The population is reproductively mature, with few of the remaining white sturgeon younger than 25 years old. The Idaho Department of Fish and Game (IDFG) estimated that 7 percent of female, and 30 percent of male white sturgeon in the Kootenai River were reproductively mature in any given year (Apperson 1992). Recent monitoring has documented an approximate 1.7:1 male to female ratio of adult fish (Paragamian et al. 1997).

The youngest white sturgeon collected in surveys since 1972 include representatives from 13 year classes (Paragamian et al. 1996, 1997). Captured fish include at least one fish hatched each year from 1972 through 1980; two fish hatched in 1983 year; and at least nine, two, and one fish produced from the 1991, 1992, and 1995 year classes, respectively. Little is known about habitats used by juvenile white sturgeon in the Kootenai River basin.

Genetic analysis indicates that Kootenai River white sturgeon are a unique stock and constitute a distinct interbreeding population (Setter and Brannon 1990). The measure of genetic variation determined for the Kootenai River population is much lower compared to white sturgeon in the lower Columbia River (Setter and Brannon 1990). Based on these comparisons, Setter and Brannon (1990) concluded "...we find adequate evidence to distinguish these fish as a separate population...." This is consistent with the geographic isolation of the population since the last glacial age.

C. Aquatic Community

Fish community associates of the Kootenai River white sturgeon include the burbot (*Lota lota*) and several native salmonids: westslope cutthroat trout

(*Oncorhynchus clarki lewisi*), interior redband and rainbow trout (*Oncorhynchus mykiss gairdneri* and *O. m. irideus*), bull trout (*Salvelinus confluentus*), kokanee, and mountain whitefish (*Prosopium williamsoni*) (Appendix A).

In general, fish populations have declined in the Kootenai River basin over the past several decades. Bull trout in the Kootenai River basin are part of the Columbia River population of bull trout listed as “threatened” in the United States under the Endangered Species Act on June 10, 1998 (63 FR 31647). Bull trout are now isolated into five subpopulations in the United States portion of the basin, with subpopulations generally with relatively low abundance. Kokanee populations have declined dramatically in the Kootenay Lake system since the 1970's. For example, kokanee runs into north Idaho tributaries of the Kootenai River numbering tens of thousands of fish as recently as the early 1980's (Partridge 1983) declined to only three fish in six of their historic spawning tributaries by 1997 (Sue Ireland, KTOI, pers. comm., 1998). Several factors are believed to have contributed to the kokanee collapse, primarily a decline in overall biological productivity due to Libby Dam construction and operations, and degraded spawning habitat. The introduction of mysid shrimp in Kootenay Lake, an efficient competitor with kokanee for food, has also contributed (Ashley and Thompson 1993). Additionally, catch rates of rainbow trout, and standing stock and growth rates of mountain whitefish in the Kootenai River have declined since the early 1980's (Paragamian 1994). The burbot population has also declined during recent decades, as indicated by an ongoing burbot population study in the Kootenai River and Kootenay Lake. The decline in burbot is not fully understood but is also thought to be partially due to the changing Kootenai River flow patterns during the winter burbot spawning period, and reduced biological productivity. Past overharvest of burbot in the Kootenai River and Kootenay Lake may also have reduced their population size (Paragamian and Whitman 1997).

D. Reasons for Decline

The significant change to the natural flows in the Kootenai River caused by flow regulation at Libby Dam is considered to be a primary reason for the Kootenai River white sturgeon's continuing lack of recruitment and declining numbers.

Beginning with the partial operation of Libby Dam in 1972 (though not fully operational until 1974), average spring peak flows in the Kootenai River have been reduced by more than 50 percent, and winter flows have increased by 300 percent compared to predam values (Figure 3). As a result of original Libby Dam operations until the initiation of experimental flows in 1992, the natural high spring flows thought to be required by white sturgeon for reproduction rarely occurred during the May to July spawning season when suitable temperature, water velocity, and photoperiod conditions would normally exist. In addition, cessation of periodic flushing flows has allowed fine sediments to build up in the Kootenai River bottom substrates. This sediment fills the spaces between riverbed cobbles, reducing fish egg survival, larval and juvenile fish security cover, and insect production.

Additionally, the elimination of side-channel slough habitats in the Kootenai River flood plain due to diking and bank stabilization to provide flood protection for agricultural land; development of Creston Valley Wildlife Management Area in British Columbia and Kootenai National Wildlife Refuge in Idaho; and lower Kootenay Lake spring maximum elevations are also a contributing factor to the white sturgeon decline. Much of the Kootenai River has been channelized and stabilized from Bonners Ferry downstream to Kootenay Lake resulting in reduced aquatic habitat diversity, altered flow conditions at potential spawning and nursery areas, and altered substrates in incubation and rearing habitats necessary for survival (Partridge 1983, Apperson and Anders, 1991).

As a consequence of altered flow patterns, average water temperatures in the Kootenai River are typically warmer (by 3 degrees Celsius; 37 degrees Fahrenheit) during the winter and colder (by 1 - 2 degrees Celsius; 34 - 36 degrees Fahrenheit) during the summer than prior to impoundment at Libby Dam (Partridge 1983). However, during large water releases and spills at Libby Dam in the spring, water temperatures in the Kootenai River may be colder than under normal nonspill spring flow conditions.

The overall biological productivity of the Kootenai River downstream of Libby Dam has been altered. Based on limnological studies of Kootenay Lake, Daley et al. (1981) concluded that the construction and operation of Libby Dam (and




Figure 3. Mean monthly Kootenai River flows at Bonners Ferry for 1928-1972 (pre-Libby Dam) and 1973-1995 (post-Libby Dam) periods.

Duncan Dam, British Columbia) "...has drastically altered the annual hydrograph and has resulted in modifications to the quality of water now entering the lake by removing nutrients, by permitting the stripping of nutrients from the water in the river downstream from Libby Dam, and altering the time at which the nutrients are supplied to the lake." Potential threats to Kootenai River white sturgeon from declining biological productivity include decreased prey abundance and food availability for some life stages of sturgeon downstream of Libby Dam, and possible reduction in the overall capacity for the Kootenai River and Kootenay Lake to sustain substantial populations of white sturgeon and other native fishes. For example, total zooplankton densities in the Kootenai River at Bonners Ferry (mean fewer than 0.1 organism/liter) are lower than in other rivers of the northwestern United States (Paragamian 1994).

Poor water quality and excessive nutrients in the upper Kootenai River were considered to be major problems for the white sturgeon and other native fishes prior to the construction and operation of Libby Dam. Graham (1981) believed that poor water quality conditions in the 1950's and 1960's, from industrial and mine development, most likely affected white sturgeon reproduction and recruitment prior to 1974. Significant improvements in Kootenai River water quality were noted by 1977, due in part to waste water control and effluent recycling measures initiated in the late 1960's. Although fertilizer processing, sewage, lead-zinc mine, and vermiculite discharges have been eliminated, many of these pollutants and contaminants persist, primarily bound in sediments.

Apperson (1992) noted detectable levels of aluminum, copper, lead, zinc, and strontium, along with polychlorinated biphenyls (PCB) and pesticides, in white sturgeon egg samples from the Kootenai River. However, other than copper, detectable levels of these compounds, e.g. polychlorinated biphenyls, organochlorides, and zinc, were lower than levels found in other Columbia River basin white sturgeon that successfully reproduce. Ultimately, the overall effects of these pollutants on sturgeon reproduction and survival are unknown. Kootenai River white sturgeon eggs have been hatched under experimental hatchery conditions using both Kootenai River water and domestic city water, however the chronic effects of heavy metals on egg hatching success and the dietary pathways of larvae and young-of-the-year white sturgeon have not been investigated.

Georgi (1993) noted that the chronic effects on wild sturgeon spawning in "chemically polluted" water and rearing over contaminated sediments, in combination with bioaccumulation of contaminants in the food chain, is possibly reducing the successful reproduction and early-age recruitment to the Kootenai River white sturgeon population.

E. Conservation Measures

At present, there are several State, Federal, Tribal, and Canadian programs and conservation efforts that may help achieve recovery objectives for the Kootenai River population of white sturgeon. These measures are described below.

1. Kootenai River management activities

The following is a brief summary of the 1991 through 1997 flow releases for Kootenai River white sturgeon. These flows, considered experimental from 1991 through 1997 and concurrent monitoring of white sturgeon, were intended to identify some factors limiting successful reproduction of Kootenai River white sturgeon and help achieve recovery.

1991: In the spring of 1991, the United States Army Corps of Engineers (USACE) and Bonneville Power Administration managed flows for white sturgeon at the request of the Idaho Department of Fish and Game. Approximately 566 cubic meters per second (m^3/s) (20,000 cubic feet per second [cfs]) were released at Libby Dam for a 2 week interval during the spawning period. The Army Corps of Engineers operations provided flows of above 991 cubic meters per second (35,000 cubic feet per second) at Bonners Ferry for 15 days with water temperatures at 14 degrees Celsius (57 degrees Fahrenheit). A peak flow of 1,521 cubic meters per second (53,700 cubic feet per second) was recorded on May 19 at Porthill, Idaho. This was accomplished without storing additional water in Koocanusa Reservoir because of above normal water conditions in the Kootenai River basin. The combination of local runoff below Libby Dam and water released to meet flood control requirements provided the range of flows (Figure 4). On July 3, 13 white sturgeon eggs were collected within 100 meters (300 feet) down river from the railroad bridge at Bonners Ferry




Figure 4. Kootenai River estimated flows and water temperature observed during April through July, 1991.

(river-kilometer 245, river-mile 153) (Apperson and Anders 1991). No larval white sturgeon were found in the Kootenai River in 1991. However, four juvenile white sturgeon aged to the 1991 year class have been found in subsequent sampling.

1992: The Bonneville Power Administration and the U.S. Army Corps of Engineers attempted to manage water releases similar to 1991 at the request of the Idaho Department of Fish and Game. However, because of the poor water year, water was not released for flood control during the white sturgeon spawning season (Figure 5). In June 1992, the Bonneville Power Administration was also requested by BC Hydro (supported by the Governor of Montana's concern for the health of the reservoir fishery) and the Army Corps of Engineers to store water in Koocanusa Reservoir for recreational purposes. As a result, flows dropped from nearly 566 to 113 cubic meters per second (20,000 to 4,000 cubic feet per second) in the Kootenai River during the critical white sturgeon spawning period. No white sturgeon eggs or larvae were found in the Kootenai River (Apperson and Wakkinen 1993).

1993: In an attempt to develop a regional prelisting recovery strategy for sturgeon that would form the basis of a conservation agreement between the U.S. Fish and Wildlife Service and various agencies, the Kootenai White Sturgeon Technical Committee (Technical Committee) was formed. The Committee comprised representatives from the U.S. Fish and Wildlife Service; Idaho Department of Fish and Game; Montana Department of Fish, Wildlife, and Parks; Kootenai Tribe of Idaho; Army Corps of Engineers; Bonneville Power Administration; and several other United States and Canadian agencies. Based upon recommendations by some Technical Committee members, the Fish and Wildlife Service requested flows of 991 cubic meters per second (35,000 cubic feet per second) for a 40-day period. The Army Corps of Engineers and Bonneville Power Administration were unable to implement the request because of operating constraints of the hydrosystem, but did store 493,413,000 cubic meters (400,000 acre-feet) of water in Koocanusa Reservoir for white sturgeon experimental flows. Water released provided 566 cubic meters per second (20,000 cubic feet per second) at Bonners Ferry from June 2 through June 16




Figure 5. Kootenai River estimated flows and water temperature observed during April through July 1992

(Figure 6). Three white sturgeon eggs (one fertilized, one dead, and one unfertilized) were collected in the Kootenai River near the US 95 Highway bridge at Bonners Ferry (river-kilometer 245, river-mile 153) when water temperatures were 12 degrees Celsius (48 degrees Fahrenheit). No larval white sturgeon were found (Marcuson 1994). To date, no 1993 year class juvenile white sturgeon have been found.

On July 7, 1993, the U.S. Fish and Wildlife Service proposed to list the Kootenai River population of white sturgeon as "endangered" under the Endangered Species Act.

1994: In July 1994, the Fish and Wildlife Service issued a formal Conference Opinion on the effects of the 1994-1998 Federal Columbia River Power System (FCRPS), concluding that the proposed operation was not likely to jeopardize the sturgeon. The action proposed by the Fish and Wildlife Service was in 3 out of 10 years to 1) maintain 425 cubic meters per second (15,000 cubic feet per second) at Bonners Ferry in May; 2) increase discharge from Libby Dam to provide 566 cubic meters per second (20,000 cubic feet per second) at Bonners Ferry for 35 days during the expected spawning season; 3) ramp down and maintain 312 cubic meters per second (11,000 cubic feet per second) for 28 days at Bonners Ferry; and 4) keep flow releases constant during May through July in years when flows were provided. This action could also benefit listed salmon species in the lower Columbia River drainage.

During the 1994 runoff period, the Bonneville Power Administration and the Army Corps of Engineers stored 1,480,000,000 cubic meters (1,200,000 acre-feet) of water behind Libby Dam as part of a flow augmentation program. This water was released to stimulate natural spawning of white sturgeon (Figure 7). Flow at Bonners Ferry was held above 425 cubic meters per second (15,000 cubic feet per second) during May and was increased to 566 cubic meters per second (20,000 cubic feet per second) on June 1 and maintained for 28 days. Flow was then decreased over 3 days to 340 cubic meters per second (12,000 cubic feet per second) by July 2, and held stable over the July 4 weekend at the request of the State of Montana to benefit recreation. Libby Dam discharge was then ramped down over 5 days to 113 cubic meters per second (4,000 cubic feet per second) by




Figure 6. Kootenai River estimated flows and water temperature observed during April through July, 1993.




Figure 7. Kootenai River estimated flows and water temperature observed during April through July, 1994.

July 11, when the 1,480,000,000 cubic meters (1,200,000 acre-feet) of stored water was exhausted. A total of 213 white sturgeon eggs were collected over 19 days beginning May 15 through June 20 near Shorty's Island (river-kilometers 228.7 - 230.9; river-miles 143 - 144) and between Myrtle and Deep Creeks (river-kilometer 237.5; river-mile 147) (Kootenai Tribe et al. 1995). No live larval white sturgeon were found in the wild during 1994, however, one newly emerged larva was found in a largescale sucker stomach in early June.

The Kootenai River population of white sturgeon was listed as endangered under the Act on September 6, 1994. In the final rule the Fish and Wildlife Service stated "that there is no recent evidence of successful spawning and survival past the egg stage" and "...existing regulations and experimental flow programs have not been effective in arresting..." the decline of the species.

1995: On December 15, 1994, the Federal Columbia River Power System action agencies submitted a supplement to the 1994-1998 Biological Assessment (B.A.) to the Fish and Wildlife Service (see previous "1994" discussion). The supplement to the Biological Assessment addressed future operation of the Federal Columbia River Power System and potential impacts upon listed species. Beginning in mid-December, the Fish and Wildlife Service, National Marine Fisheries Service (NMFS), and the action agencies (the Bonneville Power Administration, Army Corps of Engineers, and the Bureau of Reclamation [BR]) formally consulted during a series of meetings and information exchanges. The Fish and Wildlife Service and the action agencies considered how the proposal to operate the Federal Columbia River Power System as described in the Supplemental Biological Opinion could avoid jeopardy to the Kootenai River white sturgeon. To consider all viewpoints, the Fish and Wildlife Service solicited comment on the January 25, 1995, draft Biological Opinion from affected State and Tribal management agencies. On March 1, 1995, the Fish and Wildlife Service issued a final Biological Opinion addressing the effects of Federal Columbia River Power System operations in 1995 and future years on the Kootenai River white sturgeon.

The final Biological Opinion described reasonable and prudent alternatives to regulate flows at Libby Dam for 1995 to 1998. Regulation of flows must be

consistent with existing treaties and laws, e.g. the International Joint Commission and the Columbia River Treaty. Operations for 1995 were more limited than those described for 1996 to 1998 because only four of the five turbines in Libby Dam were functional.

The 1995 flow augmentation program (Figure 8) was implemented as follows: Approximately 2,467,000,000 cubic meters (2 million acre-feet) of water were stored in Koochanusa Reservoir to benefit white sturgeon. Increased flows began on April 29 to achieve 433 cubic meters per second (15,300 cubic feet per second) at Bonners Ferry on May 2. Flows ranged from 425 to 482 cubic meters per second (15,000 to 17,000 cubic feet) until May 15, when Libby Dam discharge increased to about 566 cubic meters per second (20,000 cubic feet per second) by May 16, allowing local inflow to vary Bonners Ferry flows while Libby outflow was held steady. Water temperatures remained below the optimal range for white sturgeon during most of the flow augmentation period. Bonners Ferry flows ranged from 765 to 1,076 cubic meters per second (27,000 to 38,000 cubic feet per second) during this period, which ended June 26. Flows were gradually decreased to minimum Libby Dam discharge of 113 cubic meters per second (4,000 cubic feet per second) by July 22; Bonners Ferry flow was 272 cubic meters per second (9,600 cubic feet per second). Flows were again increased on July 29, reaching about 437 cubic meters per second (16,000 cubic feet per second) by August 1, primarily to benefit salmon downstream in the Columbia River. On August 10, Kootenai River flows at Bonners Ferry reached 453 cubic meters per second (16,600 cubic feet), with very low local inflows. This second peak during the normally warm summer months departs from the natural hydrograph and can cause stranding of aquatic insects and fish eggs and larvae. Similar to 1994, 163 white sturgeon eggs were recovered only near Shorty's Island at approximately 12 river-kilometer (7.5 river-mile), downstream of Bonners Ferry, and were not recovered in the river near Bonners Ferry (Anders and Westerhof 1996). Most of the fertilized eggs were less than 60 hours old and no larvae or juvenile white sturgeon from the 1995 year class have been found through March 1996.




Figure 8. Kootenai River estimated flows and water temperature observed during April through July, 1995.

1996: Temperatures in the Kootenai River at Bonners Ferry reached an early though brief level of nearly 8 degrees Celsius (46 degrees Fahrenheit) in mid-April, and Libby Dam discharges were increased from base levels to about 650 cubic meters per second (23,000 cubic feet per second) by April 13. This level was held until about April 25, and lowland runoff complemented it, reaching peaks at Bonners Ferry of about 1,200 cubic meters per second (42,000 cubic feet per second) and 1,350 cubic meters per second (48,000 cubic feet per second) during that time (Figure 9). Lowland runoff tailed off while Libby discharge was dropped to a level of 263 cubic meters per second (9,300 cubic feet per second) by about May 1. In mid-May, lowland runoff again increased, and Libby discharges also increased in response to increasing inflows from higher elevations. A series of peaks as high as 1,400 cubic meters per second (49,500 cubic feet per second) occurred by early June at Bonners Ferry as water temperatures there exceeded 7 degrees Celsius (44 degrees Fahrenheit) and dam discharges were increased to stimulate sturgeon migration and spawning. Water temperatures reached 8 degrees Celsius (46 degrees Fahrenheit) by the end of May, and 9 degrees Celsius (48 degrees Fahrenheit) by early June. Local runoff declined starting in early June, and by the end of June was only about 300 cubic meters per second (10,600 cubic feet per second). By mid-July it was well under 100 cubic meters per second (3,500 cubic feet per second). Libby discharges were gradually dropped, but with peaks added above 700 cubic meters per second (24,700 cubic feet per second) in early and mid-July to further stimulate sturgeon reproductive activity, coinciding with temperatures of 12 degrees Celsius (54 degrees Fahrenheit), and 14 degrees Celsius (58 degrees Fahrenheit) respectively. In 1996, a total of 349 eggs were collected between June 8 and June 30. No white sturgeon larvae were collected in 1996.

1997: The Kootenai River at Bonners Ferry rose above 1,414 cubic meters per second (50,000 cubic feet per second) during 1997. Exceptionally heavy precipitation and 130 percent greater than average snow pack in the drainage raised flows at Bonners Ferry to over 1,526 cubic meters per second (54,000 cubic feet per second) during April and May (Figure 10). The peak flow for 1997 reached 1,547 cubic meters per second (54,600 cubic feet per second) on May 14. Most of the flow in April and May was local inflow. As a consequence, water management at Libby Dam was primarily for flood control at Bonners Ferry and




Figure 9. Kootenai River estimated flows and water temperature observed during April




Figure 10. Kootenai River estimated flows and water temperature observed during April through July, 1997.

the Kootenai River valley. Discharge from Libby Dam was held to only 162 to 354 cubic meters per second (5,700 to 12,500 cubic feet per second) for the entire month of April. Despite these efforts, near flood conditions still prevailed in the lower portion of the drainage because of the volume of local inflow. Test flows were initiated on June 5; flows reached 1,320 cubic meters per second (46,600 cubic feet per second) on June 6. Temperature rose from about 9.1 degrees Celsius (48 degrees Fahrenheit) on June 4 to 10.1 degrees Celsius (50.2 degrees Fahrenheit) on June 6. The first test ended when flows at Bonners Ferry were reduced slightly to 1,220 cubic meters per second (43,000 cubic feet per second) by June 10 and then increased with augmented flows from Libby Dam to produce 1,270 cubic meters per second (44,700 cubic feet per second) by June 12 at Bonners Ferry, which was the beginning of the second test. Temperature during the second flow test increased from 10.1 degrees Celsius (50.2 degrees Fahrenheit) to 11.4 degrees Celsius (52.5 degrees Fahrenheit) on June 12. Following ramp down on June 13, the temperature increased to 12.3 degrees Celsius (54.1 degrees Fahrenheit) for 3 days. Flows were gradually ramped down after the second test and were as low as 357 cubic meters per second (12,600 cubic feet per second) by the end of July. A total of 75 eggs were collected between June 5 and June 24. One larval white sturgeon was collected in the Kootenai River near Myrtle Creek at river-kilometer 236 (river-mile 145).

2. Columbia River Basin Fish and Wildlife Program

The Northwest Power Act of 1980 authorized the States of Idaho, Montana, Oregon, and Washington to create a policy-making and planning body for electrical power and the Columbia River basin's fish and wildlife resources (Northwest Power Planning Council 1987). The Northwest Power Planning Council (NPPC) was created in 1980 to develop the Columbia River Basin Fish and Wildlife Program (Program). The Program was intended to protect, mitigate, and enhance fish and wildlife resources affected by hydroelectric development in the Columbia River basin in the United States. In 1987 and 1994, the Program was amended to address several issues of concern in the Kootenai River drainage (NPPC 1987, 1994). The Bonneville Power Administration, the Army Corps of Engineers, the Bureau of Reclamation, and the Federal Energy Regulatory Commission are the Federal agencies responsible for implementing the Program.

The 1987 Program directed the Bonneville Power Administration to fund the following efforts related to the Kootenai River system:

- 1) Evaluate the effect of Libby Dam operations on reproduction and rearing of white sturgeon in the Kootenai River. Section 903(b)(1)C.
- 2) Develop operating procedures for Libby Dam to ensure that sufficient flows are provided to protect resident fish in the Kootenai River and Lake Koocanusa. Section 903(a)(5). Consult with the State of Montana if a conflict occurs between meeting minimum flows in Section 903(a)(5) and maintaining reservoir levels required by Section 903(b)(1).
- 3) Determine the impact of development and operation of the hydropower system on white sturgeon in the Columbia River basin. Section 903(e)(1).
- 4) Increase the number of rainbow trout, burbot (ling), and white sturgeon in the Kootenai River. Section 903(e)(7).
- 5) Design, construct, operate, and maintain a low-capital white sturgeon hatchery on the Kootenai Indian Reservation. Explore alternative ways to make effective use of the hatchery year-round. Section 903(g)(1)(H).
- 6) Survey the Kootenai River downstream of Bonners Ferry to the United States/Canada border to evaluate the effectiveness of the hatchery and assess the impacts of water fluctuations caused by Libby Dam on hatchery outplanting of white sturgeon in the Idaho portion of the Kootenai River. Section 903(G)(1)G.

The 1994 Program amendments called for the Bonneville Power Administration to continue to fund several of the 1987 measures for the Kootenai River drainage described above, and added several additional measures including:

- 1) Develop operating procedures for Libby Dam to ensure that sufficient flows are provided to protect resident fish. Section 10.3(B)(1).
- 2) Implement the Integrated Rule Curves (IRCs) for Koocanusa Reservoir; refine integrated rule curves to limit Koocanusa Reservoir drawdown to protect resident fish; and review State and Tribal recommendations on the biological effectiveness of the Integrated Rule Curves. Section's 10.3(B)(2,3,4).
- 3) Fund studies to evaluate the effect of Libby Dam operations on resident fish. Section 10.3(B)(5).
- 4) Design, construct, operate, and maintain mitigation projects in the Kootenai River system and Koocanusa Reservoir to supplement natural propagation of fish. Section 10.3(B)(11).
- 5) Operate and maintain a low-capital white sturgeon hatchery by the Kootenai Tribe of Idaho (KTOI). Section 10.4(B)(1).
- 6) Release water from Libby Dam to augment river discharge during the May through July sturgeon spawning period. Section 10.4(B)(3).
- 7) Restore white sturgeon and burbot populations in the Kootenai River. Section 10.6(C)(1).

3. Kootenai River white sturgeon research and monitoring

Research on white sturgeon in the Kootenai River basin by the Idaho Department of Fish and Game began in 1978 and continued through 1982. Study results indicated that white sturgeon recruitment began to decline in the mid 1960's, and that the general lack of recruitment was most pronounced after the construction of Libby Dam in 1972. White sturgeon research and monitoring in the Kootenai River basin resumed in 1988 based on the Northwest Power Planning Council's

1987 Fish and Wildlife Program (described in 2 above). These studies are funded by the Bonneville Power Administration in an effort to identify environmental factors limiting the white sturgeon population, and to recommend appropriate conservation and management actions to restore the wild white sturgeon population. The research and monitoring program has expanded in recent years with Bonneville Power Administration funding additional monitoring efforts by Montana Department of Fish, Wildlife, and Parks; Kootenai Tribe of Idaho; and British Columbia Ministry of Environment, Lands, and Parks, in addition to efforts by Idaho Department of Fish and Game. Much of the information generated from these studies was used by the Fish and Wildlife Service in the original listing determination and by the recovery team in developing this final recovery plan.

4. Kootenai Tribe of Idaho White Sturgeon Hatchery

The Kootenai Tribe of Idaho white sturgeon hatchery began as an experimental program in 1990 in response to questions concerning water quality, white sturgeon gamete viability, and feasibility of aquaculture as a component in recovery. Culture efforts first documented successful egg fertilization, incubation, egg viability, and juvenile white sturgeon survival (Apperson and Anders 1991). In 1991, 1992, 1993, and 1995, progeny from wild adult white sturgeon were successfully hatched and reared in the hatchery. The release of 305 hatchery reared age-1 and age-2 fish in 1992 and 1994 provided the first habitat use, movement, survival, and growth information for juvenile white sturgeon in the Kootenai River system. Subsequent monitoring results indicate that survival of these released fish is high and growth normal. In April and October 1997, 2,283 juvenile white sturgeon from the 1995 year class were released into the Kootenai River. Target release numbers for the conservation aquaculture program will be adjusted as more information on survival of hatchery reared juveniles becomes available.

5. Kootenai River Aquatic Investigations

Several studies authorized for the Kootenai River under the Program (as summarized in Conservation Measure #2) have been initiated or completed since

1983. These studies include:

Burbot and Rainbow Trout and Fisheries Inventory: Idaho Department of Fish and Game began the study in 1993 with the objectives to (1) identify factors that are limiting populations of burbot, rainbow trout, and other fish populations within the Kootenai River drainage in Idaho and British Columbia, and recommend management alternatives to restore the fishery to sustainable levels; and (2) determine if the burbot population is being limited by reproductive success, survival, and/or the recruitment of young burbot. Mitochondrial DNA analysis has indicated there may be two or more stocks of burbot in the Kootenai River basin (Paragamian et al. *in press*). Haplotypes from burbot collected from the Idaho and British Columbia reach of the Kootenai River were significantly different from burbot captured from two other locations within the Kootenai River drainage in Montana. A Kootenai River burbot recovery committee was formed during the spring of 1998 to devise methods and programs to restore this population.

Kootenai River Sediment and Water Quality Investigation: In 1995, the Kootenai Tribe of Idaho completed a 15-month investigation to determine if heavy metal pollutants from past mining, fertilizer production, and industrial and agricultural uses were present in the Kootenai River water column and river bed sediments. Eight sites were sampled monthly from Eureka, Montana downstream to Porthill, Idaho. Water and sediment samples were analyzed for arsenic, copper, lead, chromium, zinc, iron, mercury, selenium, and manganese. Analytical results from the water samples indicated the following pollutants violate Environmental Protection Agency aquatic criteria at several sites: mercury, lead, and selenium. Arsenic, copper, and lead were also found in river sediments. Preliminary study results concluded that at various sites, the river bottom is moderately polluted. The study has been funded for an additional 5 years to continue investigations of the biological, chemical, and limnological characteristics of the Kootenai River.

Kootenai River Ecosystem and Fishery Improvement Study: Beginning in 1995, the Kootenai Tribe of Idaho was contracted by Bonneville Power Administration to describe the existing biological community and nutrient availability in the Kootenai River. The study results will include an evaluation on the possible

effects of Libby Dam operations on the biotic community and water quality, as well as remedies for any problems identified.

Ecosystem Metabolism and Nutrient Dynamics: In 1996, Idaho State University completed a comprehensive nutrient study funded by the Bonneville Power Administration for the Kootenai River in relation to flow enhancement. Study results revealed that Lake Kootenai retained approximately 63 percent of its total phosphorus and 25 percent of its total nitrogen loading. Thus, the reservoir acts as a nutrient sink and the river downstream is nutrient deprived. Lake Kootenai does not appear to chemically stratify. Thus, selective withdrawal from areas of nutrient concentrations is not currently possible. An energy budget developed for the river basin indicated that during most sampling periods, the river was dependent upon sources of energy other than that supplied directly by within-reach autotrophic productivity. Further analysis indicated that macroinvertebrates were not energy (food) limited.

Instream Flow Incremental Methodology study: A study to determine white sturgeon habitat availability in the Kootenai River downstream of Libby Dam under various flow regimes is being conducted by the Montana Department of Fish, Wildlife, and Parks. Microhabitat investigations will be completed during 1998. Model analyses have begun and results specific to white sturgeon and associated prey organisms will be available in 1999.

Kootenai Basin Trout Genetic Analysis: Recent genetic analysis of trout species inhabiting the Kootenai River drainage indicates that interior redband trout, westslope cutthroat, and bull trout were native species in portions of the Kootenai drainage prior to development (Huston 1995). Interior redband trout still exist in the drainage, and are genetically distinct from Garry rainbow trout native to Kootenai Lake. Prior to Huston's genetic assessment, it was believed that interior redband were native only in areas downstream of Kootenai Falls (Sage et al. 1992; Behnke 1992). Populations of genetically pure redband trout were located in the Yaak River drainage and upstream of Kootenai Falls. Additional sampling is presently underway to establish the range of interior redband trout in the Kootenai River drainage upstream of Kootenai Falls.

6. Kootenay Lake Fertilization Experiments

The British Columbia Ministry of Environment, Lands, and Parks and BC Hydro are currently fertilizing the North Arm of Kootenay Lake to increase biological productivity and restore native fish populations (Ashley and Thompson 1993). This program was initiated in 1992 in response to a long-term decline in the kokanee population, especially stocks from the North Arm of Kootenay Lake. These declines raised concerns for the future of the Kootenay Lake sport fishery, dominated by the Gerrard rainbow trout. Conversely, increasing overall biological productivity in Kootenay Lake should benefit white sturgeon by increasing a potential prey base.

The project involves releasing liquid fertilizer into a 16-kilometer (10-mile) zone of the North Arm of Kootenay Lake once per week from late April through early September. The fertilizer formulation is a blend of ammonium polyphosphate (10-34-0) and urea-ammonium nitrate (28-0-0). Approximately 317 tons of 10-34-0 and 581 tons of 28-0-0 are released each year during the application period, which is the equivalent of 70 percent of preimpoundment (1949) loading levels. As of early 1997, physical limnology parameters such as temperature, dissolved oxygen, pH, Redox potential, and water clarity have not changed significantly. However, total phosphorus concentrations have increased to preimpoundment levels, which is the target for the fertilizer loadings. Additionally, algal biomass levels in the fertilized area have increased similarly. Both mysid shrimp and kokanee abundance have increased. To date, the number of kokanee spawners in two tributaries of the North Arm (Meadow Creek and Lardeau River) have ranged from a low of 300,000 in 1991 to 1.5 million in 1997.

7. Harvest Regulations

There is no legal fishing for white sturgeon within the Kootenai River drainage in either the United States or Canada (Table 1).

Table 1. Summary of historical harvest regulations for white sturgeon within the Kootenai River drainage in the United States and Canada.

Year	Idaho	Montana	British Columbia
1944	two in possession; no yearly limit; no commercial harvest		
1948	one setline; one in possession		
1949	one setline; one in possession; 76 centimeters minimum size		
1952			setlines permitted; one per day; 92 centimeters minimum size
1955	one setline; one in possession; 102 centimeters minimum size		
1957	one setline; two per year; 102 centimeters minimum size	setlines permitted for burbot only	
1960	one setline; two per year; one in possession; 92 - 183 centimeters length restriction		
1968		setline permitted for sturgeon February 15 through June 30	

Year	Idaho	Montana	British Columbia
1973		six setlines with six hooks/ line, season Feb 15 to June 30; two per year; 102 - 183 centimeters in length	
1975		no setlines permitted; two per year; 102 - 183 centimeters length restriction	
1978			100 centimeters minimum size
1979	two per year; one in possession; 92 - 183 centimeters length restriction; permit required	all fishing prohibited	
1981			one per year; 100 centimeter minimum size
1982			sturgeon declared a sport fish
1983	setlines prohibited; July 1 to December 31; one per year; 92 - 183 centimeters length restriction		
1984	catch and release only; open all year		
1989			setlines prohibited
1990			catch and release only
1994	fishing prohibited		fishing prohibited

8. Libby Reservoir Modeling

A computer model was developed by the Montana Department of Fish, Wildlife, and Parks to assess the effects of Libby Dam operations on the biota in Koocanusa Reservoir (Marotz et al. 1996). The model design was based on empirical data (field collections) from 1982 to 1995. Model components representing the physical environment and biological trophic levels were calibrated separately to assure reliable output. Model studies were used to develop Integrated Rule Curves (IRC) for Libby Dam operation. The Integrated Rule Curves contain variable reservoir drawdown and refill targets dependent on monthly inflow forecasts. Reservoir elevations and dam discharges resulting from the Integrated Rule Curves are designed to balance the many demands on Kootenai River drainage waters (including sturgeon recovery measures) with fisheries in the headwaters and salmon recovery actions in the lower Columbia River system, power production, and flood control. One aspect of the Integrated Rule Curves concept contains "tiered" water releases to simulate a natural spring runoff event to aid white sturgeon spawning and rearing. The amount of flow augmentation is proportional to water availability (drought to flood) in a given year. Water stored for later release improves annual reservoir refill probability.

9. Kootenai River Model

In 1997, through a series of workshops, an Adaptive Environmental Assessment (AEA) model for the Kootenai River was developed as part of an adaptive management process to examine the potential benefits and impacts of alternate flow regimes from Libby Dam on white sturgeon recruitment and other resources in the system. The main objective for developing the model was to provide a tool that would aid in design of an experimental management program to define management measures that would benefit white sturgeon juvenile recruitment. The discussions and data synthesis required to develop the model, and the model simulations were used to eliminate unlikely hypotheses for sturgeon recruitment decline and to eliminate policies that provided unacceptable outcomes for other resources in the system. The model consists of three main components: 1) a hydrology submodel that uses historic inflows into Libby Reservoir and tributaries, and a reservoir operation simulation (for Libby, Duncan, and Corra

Linn dams) to allow users to develop realistic discharge scenarios; 2) an aquatic production submodel that simulates turbidity, nutrient dynamics, and macroinvertebrate production in the Kootenai River; and 3) a fisheries submodel that simulates the effects of various habitat impacts related to dam operations and other watershed changes (e.g. declining nutrient loading, flood plain development) on population dynamics of white sturgeon, kokanee, burbot, rainbow and redband trout, squawfish, and other species. The model simulations summarize the tradeoffs between power economics, flood protection, and fisheries benefits, as well as tradeoffs among species associated with different flow regimes.

F. Strategy for Recovery

Recovery of Kootenai River white sturgeon is contingent upon reestablishing natural recruitment, minimizing additional loss of genetic variability to the population, and successfully mitigating biological and physical habitat changes caused by human development within the Kootenai River basin and the construction and operation of Libby Dam. This recovery plan proposes conservation actions to benefit white sturgeon within the entire Kootenai River watershed in the United States and Canada. However, the Endangered Species Act does not impose any restrictions or commitments on Canada. This recovery plan describes a strategy for improving coordination and cooperation between the United States and Canada on the operation of Libby Dam with the operation of other hydroelectric facilities within the Kootenai River basin and elsewhere in the Canadian portion of the Columbia River basin. If required for recovery, a United States - Canada binational agreement could be entered into to aid Kootenai River white sturgeon recovery, as occurred for the endangered whooping crane.

Implementation or scheduling of tasks is also based on a priority system. Priority 1 tasks are those actions that must be taken to prevent extinction or to prevent the species from declining irreversibly in the foreseeable future. Priority 2 tasks are those actions that must be taken to prevent a significant decline in species population and habitat quality, or some other significant negative impact short of extinction. Priority 3 tasks are all other actions necessary to provide for full recovery of the species. Proposed actions for native fishes have not been assigned a priority number. However, information from these actions will be useful to

evaluate how resident fish are affected by conservation actions proposed for Kootenai River white sturgeon.

Actions (or tasks) that will have the highest priority for implementation include:

Restore natural recruitment to the Kootenai River white sturgeon population (Priority 1).

Recovery will require that suitable Kootenai River ecosystem functions, including augmented seasonal Kootenai River flows, are restored to ensure habitat conditions necessary for successful white sturgeon reproduction and recruitment, i.e. survival of juveniles during their first year of life and beyond. The first stated purpose of the Endangered Species Act is, “. . . to provide a means whereby the ecosystems upon which endangered species and threatened species depend may be conserved.” The continued preservation of the sturgeon solely through artificial propagation would not be considered recovery.

Use conservation aquaculture to prevent the extinction of Kootenai River white sturgeon (Priority 1).

One recovery objective for the Kootenai River white sturgeon population is to prevent extinction by developing and implementing, for at least the next 10 years, a conservation aquaculture program, i.e. hatchery propagation. A conservation aquaculture program will include protocols on broodstock collection, gene pool preservation, broodstock mating criteria, juvenile rearing, fish health, and stocking.

Monitor the survival and recovery of the Kootenai River white sturgeon and its ecosystem (Priority 1, 2, and 3).

Concurrent with efforts to restore natural recruitment and prevent the extinction of the Kootenai River white sturgeon, further research and monitoring are necessary on life history and habitat requirements of white sturgeon and other aquatic species within the Kootenai River ecosystem.

This information is essential to understand the population dynamics of other fish species and allow resource managers to evaluate the effectiveness of conservation measures in meeting recovery goals.

Update and revise recovery plan criteria and objectives (Priority 2).

The Recovery Plan for the White Sturgeon: Kootenai River Population will be updated and revised as additional information becomes available, recovery tasks are accomplished, and as environmental conditions change.

PART II - RECOVERY

A. Recovery Objectives

The short-term recovery objectives of this recovery plan (Plan) are to a) reestablish natural recruitment to the Kootenai River population of white sturgeon and b) prevent extinction through conservation aquaculture. Proposed recovery actions include providing additional Kootenai River flows to reestablish natural recruitment and using conservation aquaculture, i.e. hatchery propagation, to prevent extinction. Due to uncertainties in egg-through-yearling survival for wild white sturgeon and the general lack of recruitment since the mid-1960's, conservation aquaculture should be used to rear juvenile white sturgeon for release into the Kootenai River, and possibly Kootenay Lake, in each of the next 10 years. The Kootenai River white sturgeon population could be considered for downlisting to threatened status in approximately 10 years if downlisting criteria described in section B. Recovery Criteria below are achieved.

The long-term objectives are to provide suitable habitat conditions and restore an appropriate age structure and effective population size to ensure a self-sustaining Kootenai River population of white sturgeon.

Recovery actions proposed in this final Plan are intended to balance white sturgeon recovery with requirements for other fish species and recreational fisheries (Executive Order 12962 of June 7, 1995) within the Kootenai River drainage. In all but the most extreme low water years, the Plan should complement conservation measures designed by the National Marine Fisheries Service to meet Snake River chinook and sockeye salmon recovery objectives downstream in the Columbia River.

B. Recovery Criteria

Criteria for reclassification or downlisting to threatened status for Kootenai River white sturgeon include:

1. Natural production of white sturgeon occurs in at least 3 different years of a 10-year period. A naturally produced year class is demonstrated through detection by standard recapture methods of at least 20 juveniles from that class reaching more than 1 year of age, and;
2. The estimated white sturgeon population is stable or increasing and juveniles reared through a conservation aquaculture program are available to be added to the wild population each year for a 10-year period. For this purpose, a year class will be represented by the equivalent of 1,000 one-year old fish from each of 6 to 12 families, i.e. 3 to 6 female parents. Each of these year classes must be large enough to produce 24 to 120 white sturgeon surviving to sexual maturity. Over the next 10 years, the number of hatchery reared juvenile fish released annually will be adjusted depending upon the mortality rate of previously released fish and the level of natural production detected. Additionally, if measures to restore natural recruitment are successful, the conservation aquaculture program may be modified. Conversely, the Fish and Wildlife Service may recommend that the conservation aquaculture program be extended beyond 10 years if adequate natural recruitment to support full protection of the existing Kootenai River white sturgeon gene pool is not clearly demonstrated, and;
3. A long-term Kootenai River Flow Strategy is developed in consultation of interested State, Federal, and Canadian agencies and the Kootenai Tribe at the end of the 10-year period based on results of ongoing conservation actions, habitat research, and fish productivity studies. This strategy should describe the environmental conditions that resulted in natural production, i.e. recruitment (as described in criterion No. 1), with emphasis on those conditions necessary to repeatedly produce recruits in future years.

Recovery or delisting will be based on providing suitable habitat conditions and restoring an effective population size and age structure capable of establishing a self-sustaining Kootenai River population of white sturgeon. Specific delisting recovery criteria will be developed as new population status, life history, biological productivity, and flow augmentation monitoring information is

collected. However, it will be approximately 25 years following approval of this recovery plan before delisting of the white sturgeon population can be considered. Twenty-five years is the approximate period for female white sturgeon added to the population during the next 10 years to reach maturity and reproduce to complete a new generation or spawning cycle.

Actions Needed to Initiate Recovery:

- o Identify and restore white sturgeon habitats necessary to sustain white sturgeon reproduction (spawning and early age recruitment) and rearing while minimizing impacts on other uses of Kootenai River basin waters, e.g. recreational facilities and the resident fishery in Koocanusa Reservoir, Kootenay Lake, and Kootenai River.
- o Develop and implement a conservation aquaculture program to prevent the extinction of Kootenai River white sturgeon. The conservation aquaculture program will include protocols on broodstock collection, gene pool preservation, propagation, juvenile rearing, fish health, and preservation stocking.
- o Work within operational guidelines for Libby Dam based upon Kootenai Integrated Rule Curves (KIRC) to balance white sturgeon recovery with requirements for other fish species and recreational fisheries within the Kootenai River drainage, and VARQ to ensure that more water is available for white sturgeon, salmon, and all species in lower water years.
- o Continue research and monitoring programs on life history, habitat requirements for all life stages, population status, and trends of the Kootenai River white sturgeon.
- o Protect Kootenai River white sturgeon and their habitats using available regulatory mechanisms, including section 7 and 10 of the Endangered Species Act, section 404 of the Clean Water Act, and the Canadian Fisheries Act.

- o Evaluate how changes in biological productivity in the Kootenai River basin affect white sturgeon and their habitats.
- o Evaluate the effects of contaminants and possible additional biological threats, i.e. predation, on Kootenai River white sturgeon and their habitats.
- o Increase public awareness of the need to protect and recover the Kootenai River white sturgeon.
- o Balance white sturgeon recovery measures with requirements for other aquatic species and recreational fisheries within the Kootenai River drainage.
- o Secure funding for implementation of recovery tasks.

Recovery of the Kootenai River population of white sturgeon will require improved coordination between United States and Canadian governmental and nongovernmental organizations. In this Plan, the Fish and Wildlife Service acknowledges numerous programs underway through local, State, Tribal, Federal, and Canadian entities to address Kootenai River basin issues. Improved interagency coordination will ensure that these, and future programs, are compatible with recovery objectives proposed for the Kootenai River white sturgeon. Additionally, a United States - Canada binational agreement could be entered into to aid Kootenai River white sturgeon recovery efforts, as occurred for the endangered whooping crane.

The Fish and Wildlife Service will use the results of ongoing research and monitoring to update and revise the plan as needed.

C. Recovery Measures Narrative

Figure 11 outlines the proposed Kootenai River white sturgeon recovery measures. Recovery tasks 11, 21, 22, 23, 24, 25, 26, 32, 41, and 42, described as follows, are short-term recovery measures essential to prevent extinction of Kootenai River white sturgeon.




Figure 11. Flow chart summarizing Kootenai River white sturgeon recovery measures.




Figure 11. Continued




Figure 11. Continued




Figure 11. Continued

1 Restore Kootenai River white sturgeon natural recruitment using flow augmentation.

Recovery of the Kootenai River white sturgeon will require providing suitable habitat conditions so that the remaining wild white sturgeon can successfully reproduce and recruit as juveniles (greater than age 1) to the population. Restoring natural recruitment to ensure a self-sustaining white sturgeon population will require implementing new operational guidelines for Libby Dam such as using tiered flows (Kootenai Integrated Rule Curves) to set aside water volumes for spring sturgeon flows and VARQ (an enhanced flood control protocol) to ensure that more water is available for white sturgeon, salmon, and all species in lower water years. The VARQ is an alternative flood control protocol developed by the U.S. Army Corps of Engineers for regulating flood control at Libby Dam, while the Kootenai Integrated Rule Curves (KIRCs) are designed to balance white sturgeon recovery with requirements for other species and recreational fisheries within the Kootenai River basin. The effects of operations at Libby extend well beyond the Kootenai basin, and flow management decisions must consider resources throughout the Columbia River basin. Factors other than flow possibly affecting white sturgeon recruitment, i.e. contaminants, predation, biological productivity, are addressed in recovery tasks # 311 through 342.

11 Adopt “adaptive” operational guidelines for Libby Dam.

Specific flow requirements for natural white sturgeon spawning and successful recruitment in the Kootenai River remain largely unknown. Until flows that contribute to successful recruitment are established, annual Kootenai River flow augmentation for white sturgeon should be based on water availability in the upper Kootenai River basin. This Plan proposes working within Libby Dam operational guidelines based upon increasing reservoir refill probability by adopting an operations model such as (KIRCs) that balances white sturgeon flow targets with Koocanusa Reservoir water levels and other aquatic resources in the Kootenai River basin, and using flood control operations like VARQ to ensure additional water is available for white sturgeon, salmon, and all species in lower water years. Under these “adaptive” operational guidelines, flow targets

will vary annually by water temperature, water volume, duration, and shape. The effects of flow and water temperature on various life stages of white sturgeon will also be monitored. This operational strategy was designed to balance resident fish concerns with power production, flood control, and Kooconusa Reservoir refill under varying water availability ranging from drought to flood conditions (Appendix C).

111 Conduct public review and approve new operational guidelines for Libby Dam.

Implementation of new reservoir operational guidelines will require improved coordination with Canadian water management entities such as the British Columbia Ministry of Environment, Lands, and Parks; Canada Department of Fisheries and Oceans; and hydro power producers such as BC Hydro. In addition, the adoption of new reservoir operational guidelines could be affected by the National Marine Fisheries Service's section 7 requirements relative to flows for listed Snake River salmon. In recognition of the need to meet the conservation requirements for sturgeon, salmon, and bull trout listed under the Endangered Species Act, the Fish and Wildlife Service and National Marine Fisheries Service will continue coordination on operations and flow augmentation programs with the goal of providing sufficient water for all listed species. Following final National Environmental Policy Act documentation and review, the North Pacific Division of the Army Corps of Engineers would issue a Record of Decision on adoption of new operational guidelines for Libby Dam.

112 Implement new operational guidelines to provide annual flow regimes to benefit white sturgeon in the Kootenai River basin.

Following completion of recovery task 111 and implementation by the Army Corps of Engineers of the new operational guidelines to manage Libby Dam operations, annual Kootenai River flow targets

will be selected based on forecasted inflow volumes (i.e. reservoir inflow expected during April 1 through August 30 in million acre-feet). White sturgeon flow targets would represent minimum flows at Bonners Ferry (i.e. Libby Dam discharge plus unregulated runoff between Libby Dam and Bonners Ferry). There would be no specific Libby Dam flow augmentation for white sturgeon in an extended drought or low water years, e.g. critical water years (less than 4.8 million acre-feet), unless increased discharges are required for emergency flood control (see Appendix C for a more complete description).

Proposed water volume released for white sturgeon will be estimated using monthly volume runoff or inflow forecasts beginning in January. The final augmentation volume will be based on the May 1 forecast (Table 2). When the forecast underestimates the actual inflow volume, minimum white sturgeon flow targets may be exceeded as excess water is released to slow the rate of reservoir refill. Overestimation of seasonal runoff may impact Koochanusa Reservoir refill by releasing water to achieve the minimum white sturgeon flow target.

Actual water releases from Libby Dam by the Army Corps of Engineers and Bonneville Power Administration during April through August will be based upon section 7 consultation with the Fish and Wildlife Service (task 118) and fine tuned through in-season management based on known in-river conditions and recommendations by several coordinating entities as described in task 113.

Table 2. "Tiered" volumes of water for sturgeon flow enhancement to be provided at Bonners Ferry according to the April-August volume runoff forecast at Libby. Actual flow releases would be shaped according to seasonal requests from the Fish and Wildlife Service and in-season management of water actually available. Volumes are in addition to the Libby minimum release of 4,000 cfs. (maf = million acre feet)

Forecast runoff volume (maf) at Libby	Sturgeon flow volume (maf) at Bonners Ferry
$0.00 \leq \text{forecast} < 4.80$	0.71
$4.80 \leq \text{forecast} < 6.00$	1.42
$6.00 \leq \text{forecast} < 6.70$	1.77
$6.70 \leq \text{forecast} < 8.10$	2.56
$8.10 \leq \text{forecast} < 8.90$	3.89
$8.90 \leq \text{forecast}$	4.77

113 Coordinate Libby Dam flow releases during April through August to achieve a suitable range of water temperature and discharge volume for successful white sturgeon recruitment.

The adoption of new reservoir operational guidelines will provide flexibility to assure that the flow augmentation for successful white sturgeon recruitment corresponds with suitable water temperatures. At Libby Dam, operators are able to release or selectively withdraw reservoir water from appropriate depths to achieve a more natural temperature regime as measured at Bonners Ferry. As appropriate water temperatures (10 to 14 degrees Celsius [50 to 57 degrees Fahrenheit]) become available at the appropriate outlet depth, and in consideration of ambient weather conditions and tributary additions downstream, Libby Dam discharge can be regulated to achieve the optimal mix of Kootenai River flow and temperature.

Annual flow management plans to manage water releases from Libby Dam during April through August will be based on coordination between the Army Corps of Engineers, Bonneville Power Administration, National Marine Fisheries Service, Fish and Wildlife Service, and other coordinating entities (e.g., State of Montana; Kootenai - Salish Nation; British Columbia Ministry of Environment, Lands, and Parks; Canada Department of Fisheries; and BC Hydro), and implemented through the Regional Forum's Technical Management Team or its successor. These entities will use a systematic approach to evaluate (task 321) how flow shaping, timing, water volume, water depth, water temperatures, prespawning flows, and substrate type may affect white sturgeon spawning behavior and recruitment. For example, the flow management plan would consider water availability in a given year and attempt to shape flows to mimic Kootenai River flows and water temperatures observed in years when some white sturgeon recruitment occurred, e.g. 1970, 1974, 1980, and 1991. White sturgeon should respond to these stimuli by forming prespawning

aggregate groups below Bonners Ferry in anticipation of moving upstream to spawn where suitable incubation, water temperature, water depths, water velocities, and substrate type exist.

Evaluating the success of an annual flow management plan will be part of task 321; with success partially defined as detecting white sturgeon eggs spawned into suitable habitats and documenting some level of natural recruitment.

At present, the Army Corps of Engineers and Montana Department of Fish, Wildlife, and Parks have an agreement to release water no closer than 16 meters (50 feet) beneath the Koocanusa Reservoir surface elevation to reduce the loss of fish (primarily kokanee) through the turbines (entrainment). Recent sampling of fish entrainment (Skaar et al. 1996) revealed that downstream losses of various fish species are severe in June, particularly when reservoir levels are low. However, to achieve temperature criteria for white sturgeon spawning in the Bonners Ferry reach, it has been necessary to withdraw surface water (upper 10 to 11 meters [30 to 35 feet]) from Koocanusa Reservoir during May and June. Methods of reducing entrainment should be pursued as part of the annual coordination to balance the effects of thermal control and flow augmentation on the reservoir fishery.

114 Investigate storing water in Koocanusa Reservoir prior to spring runoff to achieve white sturgeon flow targets.

The Montana Department of Fish, Wildlife, and Parks has shown that storing water behind Libby Dam during the winter period not only increases water availability for white sturgeon flow augmentation but also reduces impacts to the Koocanusa Reservoir fishery. By storing water for white sturgeon, reservoir elevations should remain more favorable for biological production and refill probability will be enhanced. Water releases for sturgeon then continue downstream to aid juvenile anadromous fish migration to the Pacific Ocean. Bull trout, west-slope cutthroat trout, rainbow

trout, and possibly burbot, in the Kootenai River may respond favorably to this operating strategy because the timing of releases corresponds with their life cycle requirements.

The VARQ was a flood control strategy developed by the U.S. Army Corps of Engineers while the KIRCs incorporate a flood control strategy that is compatible with flood control rule curves for Libby Dam under evaluation by the Army Corps of Engineers. The rule curves of both of these operational guidelines facilitate storing additional water prior to the spring runoff. The Army Corps of Engineers, in coordination with appropriate United States and Canada fishery agencies, should complete their analysis as it may allow for the storage of additional water available for white sturgeon flow augmentation and the minimization of impacts to Kootenai Reservoir through more frequent refill.

115 Conduct agency coordination for implementing white sturgeon flow augmentation program.

The Army Corps of Engineers, Bonneville Power Administration, Fish and Wildlife Service, National Marine Fisheries Service, First Nations, BC Hydro, appropriate States, and Canada will require specific information to plan and implement annual Kootenai River white sturgeon recruitment flow proposals. These entities should coordinate annually to ensure that regional flood control requirements will be met, adequate water volume is stored in Kootenai Reservoir, and system power needs and regional aquatic resource issues are addressed in years when white sturgeon flow augmentation will occur.

Prior to implementing the operational changes in the way water is stored and released from Libby Dam, the operating agencies should also cooperate using Columbia River Treaty protocols. Protocol V of the Treaty describes responsibilities of the entities and requires cooperation on a continuous basis to coordinate the operation of

Libby Dam with the operation of hydroelectric plants on the Kootenai River and elsewhere in Canada in accordance with the provisions of Articles XII (5), XII (6), IV (2a), and IV (2k) of the Treaty.

116 Kootenay Lake level evaluations.

One potential reason Kootenay River white sturgeon spawn in areas of apparent suboptimal conditions may be the result of implementing the 1938 International Joint Commission (IJC) Order, controlling the level of Kootenay Lake. The International Joint Commission, formed to ensure property rights are not impacted by actions of the neighboring countries, responded to a proposal to construct a hydroelectric facility at the outlet of Kootenay Lake by issuing an order that effectively controlled the surface elevation of Kootenay Lake. However, with the regulation of inflows by Libby Dam, the interpretation of the International Joint Commission order has resulted in Kootenay Lake mean maximum levels being approximately 2 meters (6.6 feet) lower since the construction of Libby dam in 1972. The Fish and Wildlife Service believes the lower maximum lake elevation may contribute to the lack of successful white sturgeon reproduction by altering river stage, flow velocity, and substrate relationships in the vicinity of sturgeon spawning habitat near Bonners Ferry. Velocities are important to spawning behavior and locations. Altered river velocities resulting from these lake elevation changes could partially explain the recent observation of white sturgeon spawning in the Kootenai River farther downstream than expected and over a sand substrate where eggs may not survive. Further discussion between appropriate Canada and United States officials should occur to determine whether lake elevation should be determined based on regulated or natural inflows. Other issues of concern as part of this evaluation include effects of International Joint Commission actions on Koocanusa Reservoir refill, and seasonal flooding effects along the Kootenai River and Kootenay Lake.

117 Evaluate alternatives to increase peak Kootenai River flows.

Examine alternatives to reliably provide peak flows in the Kootenai River at Bonners Ferry, Idaho, in the 1,100 to 1,400 cubic meters per second (40,000 to 50,000 cubic feet per second) range during the sturgeon spawning period. With the existing Libby Dam configuration two alternatives exist. The Water Resources Development Act of 1996 authorized appropriation of \$16 million to complete the installation of existing generating units 6 through 8 in Libby Dam. Since these generating units are also connected to the selective withdrawal system they could increase peak flows of temperature regulated waters by as much as 60 percent. However, flood control and public safety considerations are important to that discussion. Use of the spillway, if it were modified with flippers to reduce dissolved gas in outflows, might be an alternative to additional generating units. However, in years of high runoff, the spillway might not be available because the reservoir surface would probably be below the spillway crest elevation of 2,450 feet, for flood control purposes. Furthermore, in such situations, reservoir temperature stratification is still essentially nonexistent in early June, making warmer water difficult to obtain. This may also provide benefits to resident fish including bull trout in the Kootenai River downstream of Libby Dam if less conservative flood rule curves are adopted (Kootenai Integrated Rule Curves) and spill frequency increases.

118 Use existing authorities to conserve and restore Kootenai River white sturgeon.

Section 7(a) of the Endangered Species Act requires Federal agencies to use their authorities to carry out programs to conserve endangered and threatened species. The Fish and Wildlife Service will continue to request that the Army Corps of Engineers annually evaluate the direct and indirect effects of Libby Dam operations on

the Kootenai River white sturgeon under section 7(a).

12 Monitor effects of annual flow augmentation on Kootenai River levees and adjacent lands.

The monitoring program begun in 1995 to evaluate the physical impacts of flow augmentation on Kootenai River levees and adjacent lands downstream of Bonners Ferry should be continued by the Army Corps of Engineers and the British Columbia Ministry of Environment, Land, and Parks. The Army Corps of Engineers should identify areas where levee repairs may be necessary to protect developed areas and also identify areas where levees can be removed or left in their current state. The biological evaluation of potential impacts and benefits to resident fish and other aquatic resources will be conducted through implementing recovery tasks 32 through 562.

121 Monitor potential residential or agricultural flooding, levee erosion, pumping, and groundwater seepage resulting from flow augmentation and winter hydro peaking.

The Army Corps of Engineers' annual monitoring report for the 1995 flow augmentation program should include a description of seepage-caused inundation of agricultural lands flooded, levee erosion from peak spring flows and winter hydro peaking, and any flooding that may have resulted from white sturgeon augmentation flows. The results of this study will be useful in developing procedures and guidelines for implementing an annual levee monitoring program.

122 Identify opportunities to restore natural flood plain functions along the Kootenai River.

Based on the results of task 121, the action agencies should identify opportunities to restore natural flood plain and wetland

functions along the Kootenai River downstream of Bonners Ferry in Idaho. For example, identify landowners in flood-prone areas that may be willing to sell, lease, or assign conservation easements on portions of their land suitable for restoring natural flood plain functions. Funding may be available to implement this task through Section 206 Aquatic Ecosystem Restoration of the Water Resources Development Act of 1996.

The British Columbia Ministry of Environment, Lands, and Parks should work with the Creston Valley Wildlife Management Authority to further investigate altered Kootenai River flooding patterns to improve white sturgeon habitat.

123 Develop a public information program to explain Army Corps of Engineers past and current flood control program.

The Army Corps of Engineers should develop and distribute information on flood control operations and potential risks as part of their annual public meetings, as well as in any National Environmental Policy Act documentation of Kootenai River flow augmentation proposals.

124 Monitor effects of flow augmentation on Kootenay River levees and Kootenay Lake in British Columbia.

Proposed flow augmentation measures are designed to benefit white sturgeon reproduction primarily in the United States portion of the Kootenai River. However, physical impacts may also occur in Canada along the Kootenay River and Kootenay Lake. The British Columbia Ministry of Environment, Lands, and Parks should develop and implement a monitoring program in Canada similar to recovery task 121.

125 Assess the condition of white sturgeon spawning and incubation habitat quality, and potential substrate improvement measures.

Researchers generally agree that white sturgeon egg deposition and spawning downstream from Bonners Ferry in low velocity, silt/sand deposition areas of the Kootenai River are not currently occurring in optimal habitat for successful egg incubation, hatching, and larval rearing. An evaluation on the future use of artificial spawning and rearing substrates should be conducted. Artificial substrates have been introduced for various sturgeon species in North America, Russia, and France with varying degrees of success. These habitat projects have involved placing rock and boulder substrates in known spawning reaches of the target species.

2 Refine, implement, and evaluate a genetically sound conservation aquaculture program.

To prevent extinction of the Kootenai River white sturgeon population, a conservation aquaculture program will be implemented and evaluated for a minimum of 10 years (1999 through 2008). This program will help preserve the 10 population's remaining wild genetic variability and will begin to rebuild the natural age class structure of the wild white sturgeon population over the next 10 years. If measures to restore natural white sturgeon recruitment (described in tasks 111 to 116) are successful, the conservation aquaculture program may be adjusted before 2009. Components of this conservation aquaculture program include the following tasks:

21 The conservation aquaculture program will follow the policies and procedures of the Northwest Power Planning Council's Columbia Basin Fish and Wildlife Program and the Fish and Wildlife Service's artificial propagation policy.

All white sturgeon produced and released in the Kootenai River will be

consistent with management goals and policies. Fishery managers from the participating agencies will review existing policies and goals for consistency with the conservation aquaculture program. Additionally, they will ensure that the conservation aquaculture program is consistent with the Northwest Power Planning Council Fish and Wildlife Program and the Fish and Wildlife Service's artificial propagation policy.

211 Maintain necessary local, State, Tribal, Federal, and Canadian approval and permits for all conservation aquaculture activities.

Appropriate agencies will be properly informed of conservation aquaculture activities. Required permits for broodstock collection, transport, and release of white sturgeon in the Kootenai River system will be renewed through consultation with the Fish and Wildlife Service and appropriate State agencies. For example, a section 10 (a)(1)(A) permit authorized under the Endangered Species Act is required in order to collect, propagate, rear, and release white sturgeon.

22 Develop performance standards for the Kootenai Tribe of Idaho hatchery facilities.

Hatchery performance standards for white sturgeon are necessary to successfully spawn and rear healthy Kootenai River white sturgeon. For best results, the existing Kootenai Tribe of Idaho white sturgeon hatchery should be operated following well defined performance standards. The Kootenai Tribe of Idaho, in coordination with the Idaho Department of Fish and Game; Bonneville Power Administration; British Columbia Ministry of Environment, Lands, and Parks; Montana Department of Fish, Wildlife, and Parks; and the Fish and Wildlife Service, will develop a set of performance standards that include a description of suitable facilities, water quality standards, rearing capacities, and egg hatching/rearing protocols.

221 Maintain water quality standards for Kootenai Tribe of Idaho hatchery.

A reliable water supply with acceptable water quality is needed to ensure that healthy white sturgeon are reared in the Kootenai Tribe of Idaho hatchery. Water quality standards will be determined since the main hatchery water source is the Kootenai River. The physical characteristics of the water in the Kootenai River are variable throughout the year. Water quality factors monitored weekly by the Kootenai Tribe of Idaho include water temperature, dissolved gases, turbidity, alkalinity and hardness, nitrite, contaminants, and pathogens.

222 Upgrade Kootenai Tribe of Idaho hatchery to meet conservation aquaculture objectives.

To achieve the proposed conservation aquaculture objectives, the current Kootenai Tribe of Idaho hatchery near Bonners Ferry will require additional facility improvements and expansion. Some of the hatchery needs include additional rearing capabilities, a water sterilization system, a sediment removal system, and a supplemental oxygen system. Upgrades to the existing facility, begun in 1998, will enable the Kootenai Tribe of Idaho to remove sediment and bacteria from river water, improve water capacity, and moderately control water temperature.

223 Maintain Kootenay Trout Hatchery as secondary rearing facility.

At present, there is the risk of losing hatchery reared juvenile white sturgeon due to accidents or other unanticipated events, e.g. power outage or loss of water supply. To minimize the risk of losing one or more white sturgeon families held in the Kootenai Tribe of Idaho hatchery until fish are large enough to be marked and released, the Kootenai Tribe of Idaho will work with appropriate

Canadian officials to establish the Kootenay Trout Hatchery in Fort Steele, British Columbia as a secondary rearing or “fail-safe” facility within the Kootenai River basin.

224 Implement the conservation aquaculture program.

The Bonneville Power Administration has funded the design, development, construction, and operation of the Kootenai Tribe of Idaho hatchery since 1988 as directed by measure 10.4B.1 in the Northwest Power Planning Council Program. The hatchery successfully spawned, incubated, and reared white sturgeon in 1991, 1992, 1993, and 1995. This program is vital to the recovery of the white sturgeon population and the Bonneville Power Administration should continue to fund the Kootenai Tribe of Idaho hatchery from 1999 to 2008. The Kootenai Tribe of Idaho and Idaho Department of Fish and Game will implement the conservation aquaculture program to prevent the extinction of the Kootenai River population of white sturgeon.

23 Implement genetic preservation guidelines for broodstock collection and mating design options.

In 1993, the Bonneville Power Administration funded the development of a breeding plan for the Kootenai River white sturgeon (Kincaid 1993). The breeding plan provided a systematic approach to preserve the white sturgeon population’s genetic variability while management agencies continued work to restore Kootenai River habitat conditions necessary to reestablish natural recruitment (Appendix D).

231 Use adopted white sturgeon broodstock collection protocol.

Broodstock collected will represent the genetic variability of the population by taking representative samples with respect to run timing, size, sex, age, and other important traits to maintain long-term fitness. A broodstock collection protocol developed by the

Kootenai Tribe of Idaho and the Idaho Department of Fish and Game is summarized in Appendix E. The protocol, partially adapted from Kincaid's (1993) breeding plan, is designed to maximize collection efficiency, reproductive success, and genetic variation of broodstock while minimizing negative effects of handling stress on the natural spawning white sturgeon in the Kootenai River.

232 Collect adequate numbers of male and female broodstock to maintain the genetic variability.

Annually collect and spawn three to six females and six to nine males for broodstock (Appendix E). These fish will be held in the Kootenai Tribe of Idaho hatchery for 1 to 2 months until they are ready to be spawned. This protocol is adapted from Kincaid's (1993) breeding plan and will allow the genetic variability of the wild population to be maintained over the next 10 years.

The breeding plan incorporates a spawning matrix to minimize white sturgeon inbreeding and genetic drift. This spawning matrix is designed to maximize the diversity of genetic material passed on from artificially spawned adult white sturgeon when the hatchery-reared fish are released back into the wild population. Maximizing genetic diversity is important for the long term fitness and survival of Kootenai River white sturgeon. See Appendix D (Kincaid 1993) for more information.

233 Annually evaluate the conservation aquaculture program.

The conservation aquaculture program should be evaluated annually to ensure that the genetic variability of the Kootenai River white sturgeon population is preserved. Tissue samples from all broodstock and representative numbers of progeny are currently being archived for future electrophoretic or DNA analysis to

determine the genetic baseline for the white sturgeon population. The genetic baseline is necessary to determine if the broodstock collection protocol and spawning matrix are avoiding inbreeding and genetic drift.

24 Develop a release plan for Kootenai River white sturgeon.

A plan will be developed to govern the release of hatchery-reared fish so that conservation aquaculture objectives are met. Fish size, release time, and release locations are three factors that may affect survival of hatchery-reared sturgeon in the Kootenai River. The size of hatchery-reared sturgeon released into the Kootenai River should take into account predation and food availability to achieve maximum growth. The release plan will specify release sizes, release times, and release locations for hatchery-reared white sturgeon.

241 Evaluate appropriate production goals.

Annual production goals will range from 6,000 to 12,000 yearling white sturgeon depending on how many families are produced in any given year. This goal is designed to produce the 24 to 120 sexually mature sturgeon in each year class needed to rebuild a more natural age structure of Kootenai River white sturgeon. Based on 7 years of survey information, female and male Kootenai River white sturgeon reach sexual maturity as young as age 22 and 16 years, respectively (Vaughn L. Paragamian, IDFG, pers. comm.). White sturgeon releases should begin as soon as juvenile white sturgeon from the 1998 year class are large enough for marking, and continue for a minimum of 10 years. The production goal was developed using estimates of longevity, current survival estimates, and average age to maturity. Production goals may be altered based on the approval and future operation of a secondary backup rearing facility (see task 223).

242 Develop a fish health plan for hatchery-reared white sturgeon.

Fish health protocols will be developed to ensure that hatchery-reared white sturgeon available for release into the Kootenai River are generally healthy and disease free. Protocols will include a health inspection program for all white sturgeon life stages and prophylactic measures to prevent disease transfer in the hatchery. It is recommended that the health inspection program be administered by certified fish pathologists. These protocols will help minimize adverse impacts on the wild population and increase survival of hatchery-reared white sturgeon released into the Kootenai River basin.

243 Develop tagging protocols for hatchery-reared white sturgeon.

Permanent marking and tagging techniques are necessary to differentiate hatchery-produced white sturgeon from naturally produced white sturgeon in the Kootenai River. Protocols should use a combination of tagging methods (e.g. Passive Integrated Transponder Tags, scute removal patterns, and oxytetracycline). All fish must be permanently tagged to allow future identification by family and year class. Standardized tagging and collection methods will be developed to ensure that all appropriate information is recorded. The tagging protocol will be coordinated and approved by the Kootenai Tribe of Idaho; Idaho Department of Fish and Game; Montana Department of Fish, Wildlife, and Parks; British Columbia Ministry of Environment, Lands, and Parks; Canada Department of Fisheries and Oceans; and the U.S. Fish and Wildlife Service.

244 Develop a policy for hatchery-reared white sturgeon produced in excess of beneficial uses identified in the recovery plan.

The Kootenai Tribe of Idaho; Idaho Department of Fish and Game; Montana Department of Fish, Wildlife, and Parks; British Columbia Ministry of Environment, Lands, and Parks; Canada Department of Fisheries and Oceans; and the U.S. Fish and Wildlife Service will decide on the disposition of surplus juvenile white sturgeon. Once production goals have been met, beneficial use of surplus white sturgeon may include 1) establishment of a live gene bank or refugia population (task 245); 2) genetic analysis (mitochondrial DNA, nuclear DNA, or electrophoresis); 3) contaminant bioassays; 4) viral and bacterial research; 5) permanent marking techniques; 6) public displays and other educational purposes. Any fish remaining after all beneficial uses have been identified and addressed will be euthanized.

245 Evaluate feasibility of establishing an experimental white sturgeon population outside of the current occupied range.

When preserving any species, the probability of its persistence increases dramatically if that species exists in several populations. A nonessential experimental population of white sturgeon established somewhere in the Kootenai River basin would provide a long-term source of gene pool preservation, i.e. hatchery-reared fish, which would be available to augment the existing population if mortality rates are greater than expected or some natural catastrophe occurs. The Fish and Wildlife Service, in coordination with the affected State and Canadian entities, should evaluate the feasibility of establishing such a population, identify possible locations, e.g. Koocanusa Reservoir or Duncun Reservoir, and identify appropriate permits and disclosure documentation.

25 Release hatchery-reared white sturgeon into the Kootenai River basin.

Following completion of tasks 241 through 244, up to 1,000 juvenile white sturgeon per family will be released annually into the Kootenai River beginning in 1996. Based on the breeding plan developed by Kincaid (Appendix D and E), family releases will include the same number of juvenile sturgeon per year class to maintain the genetic variability of the Kootenai River white sturgeon population. Release times and locations will be developed to ensure optimal survival of hatchery-reared white sturgeon. Prior to release, white sturgeon will be tested for disease and visually inspected for physical deformities. Fish with obvious physical deformities will not be released and will be euthanized.

251 Adjust white sturgeon releases, as necessary, to meet objectives of the Kincaid breeding plan.

Based on implementing task 241 and using the monitoring results of recovery task 26, it may be necessary to adjust the numbers of hatchery-reared fish released in order to meet the goal of producing 4 to 10 spawning adult white sturgeon per family. Actual release numbers will be dependent upon the level of natural white sturgeon survival and recruitment detected for a given year. (Appendix D)

26 Monitor ecological interactions between hatchery-reared and wild sturgeon.

Interactions between hatchery-reared and wild white sturgeon will be monitored. A monitoring plan will be developed to ensure that hatchery white sturgeon are meeting the goals of the conservation aquaculture program. For example, survival and growth rates of released sturgeon are currently uncertain. Therefore, it is necessary to monitor released fish to determine survival and growth rates in the Kootenai River and Kootenay Lake in order to evaluate whether the Kincaid goal of producing 4 to 10 spawning adults per family spawned is being met.

261 **Determine factors limiting production (natural and hatchery) and habitat use patterns for each life history stage.**

A total of 2,588 hatchery-reared juvenile white sturgeon from 1991, 1992, 1993, and 1995 year classes have been released into the Kootenai River. Some of these fish will be captured at regular intervals to determine habitat preferences, movement, distribution, growth rate, food preferences, survival, and interactions with wild white sturgeon. This information will be used to determine habitat availability for juvenile white sturgeon, and identify additional areas to sample for wild white sturgeon spawning in the Kootenai River system.

3. **Conduct research on basic life history and monitor the level of recruitment, survival, and recovery of Kootenai River white sturgeon.**

Recovery of the Kootenai River white sturgeon can be achieved only by restoring the ecosystem upon which the fish depends. In addition to the interruption of natural spring runoff, other physical, chemical, and biological factors are believed to negatively affect the reproduction and survival of Kootenai River white sturgeon. These factors include habitat changes due to impounding water, diking, backwater habitat loss, changing levels of Kootenay Lake and the Kootenai River, altered bed-load transport rates, siltation, reduced productivity, nutrient loss, and water temperature modification. Potential biological factors include a declining effective population size, egg suffocation, lack of interstitial space, larval starvation, and predation on early life stages of white sturgeon. A better understanding of the white sturgeon life history and physical and biological factors affecting survival is necessary for developing specific recovery criteria and evaluating the success of proposed recovery measures.

31 **Describe basic life history information.**

Although much has been learned regarding the life history of Kootenai

River white sturgeon, further information regarding growth, longevity, age of maturation, migration patterns, specific spawning locations, egg, larvae and juvenile survival, and food selection is needed. This information will help document the ecological needs of the Kootenai River white sturgeon and also help to determine population viability.

311 Sample adult and juvenile white sturgeon in the Kootenai River and Kootenay Lake.

Collect biological information from captured fish including length, weight, girth, sex, pectoral fin samples for aging, and reproductive stage. This information will be useful to determine accurate age and growth rates of white sturgeon and determine environmental conditions necessary for natural reproduction and recruitment. As many as 120 sonic or radio transmitters previously attached to white sturgeon for monitoring purposes are still active or attached to free roaming fish. Most of these transmitters were attached with stainless steel wire that persists beyond the expected battery life. The Idaho Department of Fish and Game; Bonneville Power Administration; Montana Department of Fish, Wildlife, and Parks; and Kootenai Tribe of Idaho will evaluate the need to continue attaching transmitters to additional white sturgeon each year to fulfill research and monitoring needs. Only nonpermanent attachment methods should be used where feasible to ensure that transmitters remain attached only as long as necessary.

312 Collect and preserve tissue and blood samples for genetic analysis.

Tissue samples are being archived for future electrophoretic or DNA fingerprinting analysis to determine the genetic baseline for the population. This effort should be expanded basin wide to

include tissue samples from white sturgeon collected in the West Arm, North Arm, and South Arms of Kootenay Lake; and Duncun Reservoir.

313 Compile catch data to annually refine white sturgeon population size estimates in the Kootenai River basin.

Information regarding the number of juveniles and adults in the Kootenai River system is necessary to develop and prioritize short and long term recovery objectives. Catch data should be compiled and analyzed annually to determine what the natural age class structure of the population is and the effective population size are relative to recovery criteria.

314 Develop a juvenile white sturgeon year class index.

The results from annual juvenile white sturgeon sampling studies will be useful to management agencies to develop an index of annual year class strength. This method will also be useful to document the effect of flow augmentation on white sturgeon natural recruitment in meeting recovery criteria, and also detect significant differences in year-class abundance.

32 Describe Kootenai River flow requirements and Kootenay Lake elevations for natural spawning, incubation, rearing, recruitment, and survival of white sturgeon.

Specific flow requirements for natural white sturgeon spawning that result in successful recruitment are not yet well defined. However, the best available information on the relationship between Kootenai River flows and recruitment comes from collecting naturally reared recruited year classes of 1970, 1974, 1980, and possibly 1991. In these years, peak flow events at Porthill coincident with water temperature of 11 to 13 degrees Celsius (51 to 55 Fahrenheit) ranged from 708 cubic meters per second (25,000 cubic feet per second) in 1980 to 1,841 cubic meters per second

(65,000 cubic feet per second) in 1970. However, the strongest recent year class, 1974 had flow peaks of 1,416 and 1,558 cubic meters per second (50,000 and 55,000 cubic feet per second) at Porthill.

With the regulation of inflows by Libby Dam the interpretation of the Integrated Rule Curves order has resulted in Kootenay Lake mean maximum levels being more than 2 meters (6.6 feet) lower since the construction and operation of Libby Dam in 1974. We believe that lower maximum lake elevation may have contributed to the lack of successful white sturgeon reproduction in the Kootenai River by altering river stage, flow velocity, and substrate relationships in the vicinity of sturgeon spawning habitat near Bonners Ferry. Essentially, with lower Kootenay Lake levels the backwater effect of the lake is not as pronounced and therefore the white sturgeon detects suitable velocities farther downstream in the area of the sand substrates. As evidence, in 1994, 1995, and 1996, as Kootenai River peak flow and lake stage increased progressively, white sturgeon egg collections occurred increasingly farther upstream near Bonners Ferry (Paragamian et al. 1996).

Another important component of this recovery plan is to evaluate whether implementing recovery tasks 112 and 113 results in successful white sturgeon recruitment. This would entail using a systematic approach to evaluate how flow shaping, timing, water volume, water temperatures, and substrate type affect white sturgeon spawning behavior and recruitment. Also, with young-of-the-year fish produced in the system, we may begin to evaluate other factors affecting early age survival in the Kootenai River ecosystem.

321 Describe the response of spawning white sturgeon to various Kootenai River flows, water temperatures, gas supersaturation, and Kootenay Lake elevation.

Potential spawning white sturgeon will be captured and tagged with ultrasonic and radio transmitters. Both females and males will be tracked daily using telemetry gear prior to and throughout

the spawning season. Habitat used by tagged adults will be described including depth, substrate, water temperature, and mean column velocities.

Habitat use curves will be developed for white sturgeon spawning in the Kootenai River. Detailed maps of the movement and distribution of tagged sturgeon from April through September will also be developed. This information will be used to evaluate the success of proposed flow augmentation measures as described in task 11 in providing natural recruitment, and also useful in establishing habitat based recovery criteria as part of task 43.

Gas supersaturation (DGS) in the Kootenai River originating from Libby Dam may influence white sturgeon survival and riverine health. Although adult white sturgeon occupy deeper water and are less prone to gas bubble trauma, larvae and juveniles using shallow river margins and backwater sloughs may be influenced directly or indirectly (via impacts on the food supply) by elevated gas levels. Measurements of gas concentrations by the Montana Department of Fish, Wildlife, and Parks during Libby Dam spills in the 1970's revealed that saturation levels violated current Montana State water quality standards (greater than 110 percent total dissolved gas) in the Kootenai River. Supersaturated water persisted downstream beyond Kootenai Falls into the river reach inhabited by white sturgeon and their prey. This monitoring program should measure dissolved gas levels in the Kootenai River downstream of Libby Dam to assure that white sturgeon recovery is not compromised by elevated gas concentrations. Recent studies on white sturgeon larvae in the lower Columbia River revealed changes in swimming ability and increased vulnerability to predation due to gas supersaturation at sublethal exposure (Counihan et al. 1998).

This analysis will also relate the level of white sturgeon spawning and recruitment to Kootenay Lake levels. Since Libby Dam flow

regulation began, Kootenay Lake maximum spring elevations have decreased compared to pre-Libby Dam conditions. Decreased lake elevations would reduce the backwater effect of Kootenay Lake and thereby alter velocity patterns upstream in the Kootenai River. Velocities are important to spawning behavior and locations. Altered river velocities resulting from these lake elevation changes could partially explain the recent observation of white sturgeon spawning in the Kootenai River farther downstream than expected and over a sand substrate where eggs may not survive.

322 Annually measure white sturgeon spawning.

Artificial substrate mats, D-ring plankton nets, and predator fish stomachs will be used to sample eggs in the Kootenai River. Physical habitat parameters at egg collection sites will be measured including water depth, river bottom type, and mean water column velocity. Predator fish stomachs should be removed and examined for the presence of white sturgeon eggs.

Spawning can be verified by collection of eggs during the flow augmentation period. A relative index of the number of spawning episodes that occurred will be developed. Fertilized white sturgeon eggs will be analyzed to determine developmental stage. Combined with water temperature during the incubation period, this information will be used to back-calculate time of spawning and associated physical habitat parameters.

323 Measure white sturgeon larvae, fry, and juvenile abundance and distribution in the Kootenai River and Kootenay Lake annually.

Year class abundance can be determined for most types of fish during the first year of life. As yet there are no reliable techniques for determining year-class abundance of young-of-the-year or age 1 white sturgeon in the Kootenai River basin. The monitoring

program (described under task 3) will continue to use and evaluate a variety of traps, trawls, and nets to reliably sample white sturgeon larvae and fry in the Kootenai River system. Abundance estimates will be calculated annually for white sturgeon larvae and fry in the Kootenai River, along with potential larval and young-of-the-year rearing habitat. These data will provide further insight into locations of spawning and rearing habitat and fish movements.

Juvenile white sturgeon abundance and distribution will be monitored with small mesh gill nets in the Kootenai River and Kootenay Lake. Relative abundance estimates will be calculated for juvenile fish using a sampling design based on location, time of year, gill-net sampling effort, and total catch. Potential juvenile rearing habitat will also be identified. Habitat use curves will then be prepared and compared to available aquatic habitat through the use of In-stream Flow Incremental Methodology (IFIM) (task 324). Knowledge of critical life-cycle requirements will be used to evaluate and direct habitat enhancement efforts.

324 Quantify sturgeon spawning/incubation habitat and early rearing habitat using In-stream Flow Incremental Methodology.

Habitat use data developed in tasks 321 through 323 will be used in the In-stream Flow Incremental Methodology model to quantify and locate spawning habitat and early rearing habitat in the Kootenai River system at different river discharge levels. This information will be used to evaluate the response of white sturgeon to habitat available during various flow regimes.

33 Conduct investigations on biological productivity in the Kootenai River.

Koocanusa Reservoir currently acts as a nutrient sink and thus limits the primary and secondary productivity of the Kootenai River downstream of

Libby Dam. Changes in nutrient availability affect the food chain for the fish community, the prey base for many species including white sturgeon, growth rates, and possibly survival of larval fish.

331 Assess the necessity of increasing nutrients in the Kootenai River.

Similar to the Kootenay Lake fertilization project previously described in Part I - Conservation measure 5, artificial additions of phosphorus and nitrogen may be a potential means of restoring primary and secondary productivity in the Kootenai River. All existing information regarding stream fertilization should be compiled and evaluated. Following this evaluation, a program describing potential nutrient dynamics and possible benefits to Kootenai River white sturgeon recovery from stream fertilization should be developed in cooperation with appropriate Canada, Montana, Idaho, and Indian Tribes. Improved primary and secondary productivity in the Kootenai River basin will also benefit other fish species, e.g. bull trout, rainbow trout, kokanee, burbot, and mountain whitefish.

332 Use computer modeling to refine and analyze recovery tasks.

In 1997, through a series of workshops, an Adaptive Environmental Assessment (AEA) model for the Kootenai River was developed as part of an adaptive management process to examine the potential benefits and impacts of alternate Kootenai River flow regimes on white sturgeon recruitment and other resources in the system. The main objective was to provide a tool that would aid in design of an experimental management program to define a flow regime that would benefit white sturgeon juvenile recruitment. The model simulations summarize the tradeoffs between power economics, flood protection, and fisheries benefits, as well as tradeoffs among species, associated with different flow regimes. The model will be

used to evaluate the effectiveness of recovery tasks presented in this plan.

34 Evaluate the effects of contaminants on white sturgeon.

The Bonneville Power Administration funded a water and sediment quality study of the Kootenai River in the United States from Eureka, Montana downstream to Porthill at the United States/Canada border. However, lethal and sublethal effects of water and sediment chemical constituents on early life stages of white sturgeon still need to be determined.

341 Compile existing information on contaminants in the Kootenai River.

Use available information found in recent studies completed by the Kootenai Tribe of Idaho and Idaho State University to determine the presence and concentrations of contaminants including metals, organics, and inorganics in the water, sediment, and biota in the Kootenai River.

342 Conduct contaminant bioassays to evaluate the effects of selected chemicals on white sturgeon.

Laboratory studies of effects of heavy metals and other contaminants on white sturgeon eggs, larvae, and juveniles should be initiated. Existing protocols should be used where applicable; where no protocols exist, they should be developed with the cooperation of the Environmental Protection Agency.

4 Implement conservation and recovery of Kootenai River white sturgeon.

Recovery of Kootenai River white sturgeon is dependent upon regional coordination and adequate funding to implement conservation measures proposed in this plan.

41 Develop a Participation Plan to support implementation of the Kootenai River white sturgeon recovery plan.

Implementation of this recovery plan for Kootenai River white sturgeon will be accomplished only through interagency cooperation and participation leading to the timely recovery of the species while minimizing regional social and economic impacts. To meet these objectives, the Fish and Wildlife Service on July 1, 1994 issued new policy to develop a public Participation Plan for implementing recovery actions. Participation Plans are intended to ensure that a feasible recovery strategy involves and addresses the concerns of affected interest groups while providing realistic and timely recovery of the species. In the case of the Kootenai River white sturgeon, a Participation Plan would be developed by most of the agencies represented on the recovery team, and could include summaries of annual work plans for Kootenai River monitoring, research, and hatchery projects and section 7 consultations.

42 Identify funding required to achieve Kootenai River white sturgeon recovery.

Existing budgets of participating and responsible parties are not capable of funding all recovery tasks identified in this final plan. The recovery team should be retained to identify various funding strategies, including congressional appropriations, water-use fees, Federal mitigation programs, and binational agreements that may be useful in implementing white sturgeon recovery efforts.

421 Recommend research and management regarding ecosystem and fishery improvement measures for the Kootenai River basin.

As new information is developed and recovery actions are implemented, the recovery team should meet to address “new” research and management needs concurrent with white sturgeon

recovery activities. The Fish and Wildlife Service anticipates that new questions and data needs will arise as white sturgeon recovery implementation occurs. The recovery team would meet to develop specific proposals to address these data gaps and recommend possible funding sources.

43 Reevaluate downlisting and/or delisting criteria as needed.

As initial recovery measures (see tasks 1-342) are accomplished and/or additional information regarding the ecology of Kootenai River white sturgeon becomes available, specific delisting criteria will be established.

44 Determine the indirect recovery costs of foregone power generation.

Implementing the many conservation actions proposed in this recovery plan may create additional economic impacts that are not normally considered a true “cost” of recovery. These impacts include foregone power generation opportunities, flood control impacts, and resident fish impacts.

The Army Corps of Engineers, Bonneville Power Administration, and BC Hydro should conduct an economic analysis of proposed white sturgeon recovery actions in terms of foregone power generation and remedial flood control requirements. This analysis should determine if the current “base economic assumptions” regarding lost power revenues are valid. The analysis should also consider alternative regional power marketing strategies to reduce revenue impacts and identify innovative measures to reduce potential flood control costs.

45 Increase public awareness of the need to protect Kootenai River white sturgeon.

Increase public awareness of the need to protect Kootenai River white sturgeon and their habitat (or ecosystem). Specific tasks to accomplish this might include periodic news releases, brochures, interactive presentations,

in-school presentations by recovery team members, and possibly television documentaries.

5. Monitor the status of native fishes in the Kootenai River drainage.

The Kootenai River basin once provided important recreational, consumptive, and native subsistence fisheries. In addition to white sturgeon, residents and nonresidents fished for kokanee, burbot, rainbow trout, westslope cutthroat trout, bull trout, and mountain whitefish. All of these fisheries have declined dramatically over the past several decades. For example, a recent creel survey by the Idaho Department of Fish and Game revealed that fishing effort in the Idaho portion of the Kootenai River is the lowest of all waters surveyed in northern Idaho (Vaughn L. Paragamian, IDFG, pers. comm., 1996). Conversely, the abundance of nongame fish (e.g. suckers, northern squawfish) is three times higher than prior to the construction and operation of Libby Dam. Restoration of recreational fisheries is important to anglers and the regional economy.

Studies on the status of native fish in the Kootenai River basin were first authorized by the Northwest Power Planning Council in 1983. Although many of these studies continue, additional information is still needed on the status and important habitats required by several of the native and recreationally important fish species, including bull trout, kokanee, rainbow trout, burbot, and mountain whitefish. This information will also be useful to evaluate how resident fish are affected by conservation actions for Kootenai River white sturgeon.

51 Conduct studies on kokanee downstream from Libby Dam.

The Montana Department of Fish, Wildlife, and Parks; Idaho Department of Fish and Game; Kootenai Tribe of Idaho; and British Columbia Ministry of Environment, Lands, and Parks should continue annual monitoring to determine if kokanee entrained through Libby Dam during white sturgeon and salmon flow augmentation survive and contribute to downstream regional fisheries. Annual population estimates of kokanee would also be useful in determining whether increasing kokanee populations observed in recent years are affected by nutrient availability in the Kootenai River and Kootenay Lake.

511 Collect abundance, distribution, and reproduction data for kokanee in the lower Kootenai River tributaries.

Annual kokanee spawning population estimates will be determined. Information will be used to provide recommendations for improving kokanee spawning habitat and reintroducing kokanee in the Kootenai River tributaries. Additionally, the Idaho Department of Fish and Game; British Columbia Ministry of Environment, Lands, and Parks; and Kootenai Tribe of Idaho should evaluate opportunities to enhance spawning habitats in the Yaak River and Lake Creek.

52 Determine the status and distribution of burbot in the Kootenai River downstream of Kootenai Falls and Kootenay Lake.

Burbot are currently classified as a State threatened species by the Idaho Department of Fish and Game. The commercial and sport harvest of burbot prior to 1974 was estimated as high as 25,000 kilograms (55,000 pounds) in some years. This was primarily a winter fishery with few burbot caught in the spring and fall. Since that time, the burbot fishery in the Kootenai River basin has collapsed. There has been scant evidence of reproduction, only one juvenile burbot and no larvae have been captured in recent years. Sonic telemetry studies and recaptures reveal that the Goat River is the only known spawning location in the lower Kootenai River drainage (Paragamian et al 1997).

521 Determine distribution and life history of burbot.

The study begun in 1993 to identify distribution, life history, and factors limiting populations of burbot within the Kootenai River drainage should continue to be funded by the Bonneville Power Administration. All burbot captured will be tagged, and population estimates will be conducted annually to monitor population trends.

522 **Determine secondary impacts from the flow augmentation program on burbot.**

Recent research efforts by the Idaho Department of Fish and Game suggests that high, fluctuating Kootenai River flows during the winter affect winter migrations of burbot and possibly impact reproduction. Information garnered from implementing task 11 and completing task 521 should be used to evaluate how the proposed flow augmentation program will impact burbot recruitment. Preliminary study results indicate that burbot migrations during the spawning season may be effected by Libby Dam outflows during the winter for power production and flood control. Flow tests should be conducted to determine the maximum tolerable discharge and duration to allow burbot migration. This information is important because burbot in Idaho and British Columbia are genetically distinct from burbot in the Montana reach of the Kootenai River. Some believe this stock may be at greater risk of extinction than sturgeon.

53 **Identify factors limiting rainbow trout survival and/or recruitment in the Kootenai River basin.**

Rainbow trout spawning activity should be monitored to evaluate egg desiccation and/or redd scouring impacts in the Kootenai River from the white sturgeon flow augmentation program.

531 **Determine the status, distribution, and habitat use of rainbow trout in the Kootenai River from Libby Dam to Bonners Ferry.**

The Idaho Department of Fish and Game and the Montana Department of Fish, Wildlife, and Parks should further investigate the status and distribution of rainbow trout, including native Gerrard and interior redband, in the Kootenai River downstream of Libby Dam. Habitat use will be determined for fry, juvenile, and adult

rainbow trout using scuba and snorkeling in the Kootenai River. This information will be useful to evaluate the effects of white sturgeon flow augmentation on rainbow trout.

54 Develop information on bull trout in the Kootenai River basin.

On June 10, 1998, the Columbia River population of bull trout was listed as a “threatened” species (63 FR 31647) under the Endangered Species Act. Additional information is needed on life history requirements, distribution, and factors regulating bull trout subpopulations within the Kootenai River drainage.

541 Determine distribution and status of bull trout in tributaries of the Kootenai River.

Bull trout are known from the Kootenai River, Koocanusa Reservoir, Kootenay Lake, and several tributaries within the Kootenai River basin. Bull trout are currently isolated into five subpopulations in the United States portion of the basin, with subpopulations generally stable with relatively low abundance. Monitoring by Idaho Department of Fish and Game; Montana Department of Fish, Wildlife, and Parks; British Columbia Ministry of Environment, Lands, and Parks; Canada Department of Fisheries and Oceans; and the Kootenai Tribe of Idaho will better describe the distribution, abundance, and habitat availability for bull trout. For example, bull trout surveys, including redd counts, should be conducted for all Montana streams where bull trout have previously been found, including Quartz, O'Brien, Libby, and Pipe Creeks and the Fisher River.

542 Identify additional conservation measures to protect bull trout in the Kootenai River drainage in Montana, Idaho, and British Columbia.

The Idaho Department of Fish and Game; Montana Department of Fish, Wildlife, and Parks; British Columbia Ministry of Environment, Lands, and Parks; Canada Department of Fisheries and Oceans; and the Kootenai Tribe of Idaho, using information garnered from task 541, should identify additional conservation measures necessary to maintain bull trout within the Kootenai River basin. Additionally, these agencies should evaluate whether recovery measures proposed for white sturgeon impact bull trout.

55 Conduct research on mountain whitefish in the Kootenai River from Libby Dam downstream to Bonners Ferry.

Habitat use will be determined for fry, juvenile, and adult mountain whitefish using SCUBA and snorkeling. If possible, separate use data will be obtained for winter, summer, and spawning habitat. Microhabitat measurements (e.g. depth, velocity, substrate, and cover) will be taken at locations where fish are encountered.

551 Determine the secondary effects of proposed white sturgeon flow augmentation on mountain whitefish.

A secondary effect of the white sturgeon flow augmentation program would be less water available during the winter when mountain whitefish spawn. In order to meet normally high, daily power demands during the winter, Libby Dam discharge fluctuations could possibly dewater and kill incubating mountain whitefish eggs. The Montana Department of Fish, Wildlife, and Parks and the Idaho Department of Fish and Game should monitor these potential impacts.

56 Evaluate impacts of flow augmentation on resident fish in Canada and the United States.

Flow augmentation proposals to benefit white sturgeon and salmon will result in water spill at Canadian Kootenay River dams. Additional

monitoring is needed to evaluate the potential fisheries impacts on the Duncan, Arrow, and Koocanusa systems due to proposed recovery measures.

561 Effects of flow augmentation on total gas pressure in the Kootenay River and Columbia River downstream of Kootenay Lake.

Flow augmentation proposals to benefit white sturgeon will result in water spill at Canadian Kootenay River dams. This will increase total gas pressure levels to possibly lethal levels for some fish downstream of Brilliant Dam. Columbia Power Corporation, the Canada Department of Fisheries and Oceans, The British Columbia Ministry of Environment, Lands, and Parks, and Environment Canada should monitor these impacts, and consideration should be given to increasing hydroelectric capacity or using other gas reduction technology at Brilliant Dam as a means to mitigate these resident fish impacts.

562 Effects of flow augmentation on Kootenay Lake and on the Duncan and Arrow Reservoirs/Columbia River systems.

Potential fisheries impacts on the Duncan and Arrow reservoir systems due to white sturgeon flow augmentation from Libby Dam include 1) fluctuating flow releases from Duncan Dam during bull trout spawning migrations. This may affect bull trout movement and general spawning behavior; 2) decreased flow releases from Keenleyside Dam during rainbow trout spawning and rearing periods. This may reduce available spawning habitat and changes in temperature regimes due to flow changes, which may result in changes in incubation times; 3) decreased flow releases from Keenleyside may negatively effect staging and spawning of Columbia River white sturgeon; and 4) August releases from Libby Dam passing through Kootenay Lake and/or Arrow reservoir may flush nutrients and forage organisms from upper strata waters affecting overall biological productivity.

563 Evaluate effects of hydro peaking on native fish throughout the Kootenai River downstream of Libby Dam.

Daily load following and power peaking at Libby Dam may increase flows by fivefold in a few hours. These types of flows have altered the Kootenai River in the reach downstream from Bonners Ferry to Kootenay Lake to the extent that the river rarely freezes during the winter. These practices may particularly be impacting bull trout in the vicinity of Libby Dam in Montana and burbot in Idaho. Evaluations as part of tasks 51, 52, 53, 54, and 55 should include the effects of load following and power peaking.

564 Evaluate measures to reduce risk to native and resident sport fish below Libby Dam.

Demand for refill at Libby Dam for salmon recovery efforts, white sturgeon recovery, and sport fishing interests may lead to less conservative flood rule curves at Libby Dam in the 85- to 100-year protection range proposed in the original project justification. This would result in increasing the risk of spill and injury to bull trout and other native or resident sport fish since the frequency of “unregulated” spill will increase. The Army Corps of Engineers; Bonneville Power Administration; Montana Department of Fish, Wildlife, and Parks; and the Fish and Wildlife Service should evaluate “flip lips” and other structures that could minimize fish injuries and gas supersaturation downstream of Libby Dam.

6 Assess the overall success of implementation of the recovery plan and revise accordingly.

This plan should be updated on a 5-year basis as recovery tasks are accomplished, or revised as environmental conditions change and/or monitoring results or additional information becomes available.

The recovery team should meet annually to review annual monitoring reports and summaries and make recommendations to the Fish and Wildlife Service to revise the Plan.

REFERENCES CITED

- Anders, P.J. and R. Westerhof. 1996. Natural spawning of white sturgeon (*Acipenser transmontanus*) in the Kootenai River, Idaho, 1995. Preliminary Report of Research FY 95. Kootenai Tribe of Idaho and Bonneville Power Administration. 14 pp.
- Apperson, K.A. 1992. Kootenai River white sturgeon investigations and experimental culture. Annual Progress Report FY 1991. Idaho Department of Fish and Game and the Bonneville Power Administration. Contract No. DE-AI79-88BP93497; Project No. 88-65. Portland, Oregon.
- Apperson, K.A. and P.J. Anders. 1991. Kootenai River white sturgeon investigations and experimental culture. Annual Progress Report FY 1990. Idaho Department of Fish and Game and the Bonneville Power Administration. Contract No. DE-AI79-88BP93497; Project No. 88-65. Portland, Oregon. 67 pp.
- Apperson, K.A. and V.D. Wakkinen. 1993. Kootenai River white sturgeon investigations and experimental culture. Annual Progress Report FY 1992. Idaho Department of Fish and Game and the Bonneville Power Administration. Contract No. DE-AI79-88BP93497; Project No. 88-65. Portland, Oregon. 37 pp.
- Ashley, K. and L. Thompson. 1993. Kootenay Lake fertilization experiment - year 1 (1992/1993) report. Fisheries Branch, Ministry of Environment, Lands and Parks, University of British Columbia. Fisheries Project Report No. RD 32. 52 pp.
- Bayley, P.B. 1995. Understanding large river floodplain ecosystems. *Bioscience*. Vol. 45, No. 3. Pp 153-158.
- Behnke, R.J. 1992. Native trout of Western North America. American Fisheries Society. Monograph #6. 261 pp.
- Brannon, E.L., C.L. Melby, and S.D. Brewer. 1984. Columbia River white sturgeon enhancement. Final Report to Bonneville Power Administration. Contract No. DEAI79-84BP18952; Project No. 83-316. Portland, Oregon. 43 pp.

- Counihan, T.D., A. J. Miller, M.G. Mesa, and M. J. Parsley. 1998. The effects of dissolved supersaturation on white sturgeon larvae. *Transactions of the American Fisheries Society*. 127: 316-322.
- Daley, R.J., E.C. Karmach, C.B.J. Gray, C.H. Pharo, S. Jasper, and R.C. Wiegand. 1981. The effects of upstream impoundments on the limnology of Kootenay Lake, B.C. Scientific Service No. 117. National Water Research Institute, Inland Waters Directorate. Vancouver, British Columbia, 1981. 96 pp.
- Georgi, A. 1993. The status of Kootenai River white sturgeon. Report prepared for the Pacific Northwest Utilities Conference Committee. September 1993. 58 pp plus appendices.
- Graham, P.J. 1981. Status of white sturgeon in the Kootenai River. Montana Department of Fish, Wildlife and Parks. Kalispell, Montana. 26 pp.
- Junk, W.J., P.B. Bayley, and R.E. Sparks. 1989. The flood pulse concept in river-floodplain systems. *Canadian Special Publication, Fish Aquatic Sciences*. 106: 110-127.
- Huston, J.E. 1995. A report on Kootenai River drainage native species search, 1994. A report to the U.S. Fish and Wildlife Service. Montana Fish, Wildlife & Parks. 21 pp., attached GIS map.
- Kincaid, H.L. 1993. Breeding plan to preserve the genetic variability of the Kootenai River white sturgeon. Report prepared for the Bonneville Power Administration. Contract No. DE-AI79-93B002886. Portland, Oregon.
- Lindsay, R.A. 1995. Kootenay Lake white sturgeon investigations. Progress Report prepared for the IDFG. September, 1995. 8 pp.
- Marcuson, P. 1994. Kootenai River white sturgeon investigations. Annual Progress Report FY 1993. Idaho Department of Fish and Game and the Bonneville Power Administration. Contract No. DE-AI79-88BP93497; Project No. 88-65. Portland, Oregon. 67 pp.
- Marotz, B. L., D. Gustafson, C. Althen, and B. Lonon. 1996. Model development to establish integrated operational rule curves for Hungry Horse and Libby Reservoirs, Montana. Report prepared for the Bonneville Power Administration. Portland, Oregon. 110 pp.
- Northcote, T.C. 1973. Some impacts of man on Kootenay Lake and its salmonids. Great Lakes Fishery Commission, Technical Report, No. 2.

- Northwest Power Planning Council. 1987. Columbia River basin fish and wildlife program. Portland, Oregon.
- Northwest Power Planning Council. 1994. Columbia River basin fish and wildlife program.. Publication 94-55. Northwest Power Planning Council. Portland, Oregon.
- Pacific States Marine Fisheries Commission. 1992. (Hanson, D.L., editor). White sturgeon management framework plan. 200 pp.
- Paragamian, V.L. 1994. Kootenai River Fisheries Investigations: Stock status of burbot and rainbow trout and fisheries inventory. Annual Report 1993. Idaho Department of Fish and Game and Bonneville Power Administration. Contract No. DE-B179-88BP93497. 28 pp.
- Paragamian, V.L. and V. Whitman. 1997. Kootenai River fisheries investigations: stock status of burbot. Annual Report 1997. Idaho Department of Fish and Game and Bonneville Power Administration. Contract No. DE-B179-88BP93497.
- Paragamian, V.L., G. Kruse, and V. Wakkinen. 1996. Kootenai River Fisheries white sturgeon spawning and recruitment evaluation.. Annual Report 1996. Idaho Department of Fish and Game and Bonneville Power Administration. Contract No. DE-B179-88BP93497.
- Paragamian, V.L., G. Kruse, and V. Wakkinen. 1997. Kootenai River Fisheries white sturgeon spawning and recruitment evaluation.. Annual Report 1997. Idaho Department of Fish and Game and Bonneville Power Administration. Contract No. DE-B179-88BP93497.
- Paragamian, V. L., M. Powell, and J.C. Faler. In press. Mitochondrial DNA analysis of burbot Lota lota stocks in the Kootenai River basin of British Columbia, Montana and Idaho. Transactions of the American Fisheries Society.
- Partridge, F. 1983. Kootenai River fisheries investigations in Idaho. Idaho Department of Fish and Game, Boise, Idaho.
- Sage, G.K., R.F. Leary, and F.W. Allendorf. 1992. Genetic analysis of 45 trout populations in the Yaak River drainage, Montana. Wild Trout and Salmon Genetics Laboratory Report, 92/3, Division of Biological Sciences, University of Montana, Missoula, Montana.

Scott, W.B. and E.J. Crossman. 1973. Freshwater fishes of Canada, Bulletin 184. Fisheries Research Board Canada. Ottawa, Canada.

Setter, A. and E. Brannon. 1990. Report on Kootenai River white sturgeon electrophoretic studies - 1989. Report for Idaho Department of Fish and Game. Aquaculture Extension, University of Idaho. Pp. 43-50 In: Apperson, K.A., editor. Kootenai River white sturgeon investigations and experimental culture. Annual Progress Report FY 1989. Idaho Department of Fish and Game and the Bonneville Power Administration. Contract No. DE-AI79-88BP93497; Project No. 88-65. Portland, Oregon.

Skaar, D., J. DeShazar, L. Garrow, T. Ostrowski, and B. Thornburg. 1996. Quantification of Libby Reservoir levels needed to maintain or enhance reservoir fisheries: investigations of fish entrainment through Libby Dam, 1990-1994. Annual Progress Report. Montana Fish, Wildlife and Parks and the Bonneville Power Administration. Contract No. DE-AI79-83-BP12660; Project No. 83-467. Portland, Oregon.

Simpson, J. and R. Wallace. 1982. Fishes of Idaho. Northwest Naturalist Book, Moscow, Idaho. 238 pp.

Sparks, R.E. 1995. Need for ecosystem management of large rivers and their floodplains. Bioscience. Vol. 45, No. 3: 168-182.

PART III - IMPLEMENTATION SCHEDULE

The Implementation Schedule that follows describes recovery task priorities, task numbers, task descriptions, duration of tasks, potential or participating responsible parties, and lastly, estimated costs, if available. These tasks, when accomplished, will contribute to recovery of the Kootenai River population of white sturgeon as discussed in Part II of this Plan.

Parties with authority, responsibility, or expressed interest to implement a specific recovery task are identified in the Implementation Schedule. Listing a responsible party does not imply that prior approval has been given or require that party to participate or expend any funds. However, willing participants will benefit by demonstrating that their budget submission or funding request is for a recovery task identified in an approved recovery plan, and is therefore part of a coordinated recovery effort to recover the Kootenai River population of white sturgeon. In addition, section 7(a)(1) of the Endangered Species Act directs all Federal agencies to use their authorities to further the purposes of the Act by implementing programs for the conservation of threatened or endangered species.

Other physical and economic impacts from recovery

Implementing the many conservation actions proposed in this recovery plan will create additional economic or environmental impacts, and also associated benefits, not normally considered in estimating the “costs” of recovery. Economic and environmental impacts include foregone power generation opportunities, flood control impacts, and resident fish impacts.

- o Flood control impacts can include agricultural and residential flooding, groundwater seepage, and pumping costs. For example, crop losses ranging from 30 to 100 percent of a total 650 acres on 12 farms in the United States portion of the Kootenai Valley were attributed to the 1995 white sturgeon/salmon recovery flows (Dave Wattenburger, Boundary County Extension, *in litt.* 1996). The value of crop losses has not been estimated to date. Some farmlands were inundated, others were yellowed through soil saturation, and other lands were inaccessible during the growing season for

weed control activities. Irrigation drainage pumping costs for the period May 1 through July 15, 1995 were estimated at \$19,325 in the same United States portion of the Kootenai Valley. This cost will be adjusted downward when baseline pumping costs for ongoing average pumping needs are provided.

- o Nontargeted Fish impacts. Flow augmentation proposals to benefit white sturgeon and salmon will result in water spill at Kootenay River dams, in Canada. This will increase total gas pressure levels to possibly lethal levels for some fish downstream of Brilliant Dam. Impacts should be monitored and consideration should be given to increasing hydroelectric capacity or using other gas reduction technology at Brilliant Dam as a means to mitigate these impacts on resident fish.

Potential fisheries impacts on the Duncan and Arrow Reservoirs and Columbia River systems due to white sturgeon flow augmentation from Libby Dam include 1) fluctuating water releases from Duncan Dam during bull trout spawning migrations, which may affect bull trout movement and general spawning behavior; 2) decreased water releases from Keenleyside Dam during rainbow trout spawning and rearing periods, which may reduce available spawning habitat and changes in temperature regimes due to flow changes may result in changes in incubation times; and 3) decreased water releases from Keenleyside, which would negatively effect staging and spawning of Columbia River white sturgeon.

Associated benefits include the partial restoration of a more natural Kootenai River hydrograph and flood plain function that benefit resident fish and wildlife. Periodic flushing flows would cleanse Kootenai River gravels and improve insect production. Improving the aquatic ecosystem health leading to improved regional fisheries will provide secondary economic benefits to local communities. Such benefits go beyond the “benefits” typically considered in recovery actions. Conversely, failure to implement proposed recovery actions would have hidden environmental costs that are typically not considered in cost/benefit analysis.

Following are definitions to column headings and keys to abbreviations and acronyms used in the Implementation Schedule:

Priority No.: All priority 1 tasks are listed first, followed by priority 2 and priority 3 tasks.

Priority 1: Actions that must be taken to prevent extinction or to prevent the species from declining irreversibly in the foreseeable future.

Priority 2: Actions that must be taken to prevent a significant decline in species population or habitat quality, or some other significant negative impact short of extinction.

Priority 3: All other actions necessary to provide for full recovery (or reclassification) of the species.

Task Number and Task Description: Recovery tasks as numbered in the recovery outline. Refer to the Narrative for task descriptions.

Task Duration: Expected number of years to complete the corresponding task. Study designs can incorporate more than one task, which when combined, can reduce the time needed for task completion.

Responsible or Participating Party: Federal, State, Tribal, or Canadian government agencies, nongovernment organizations, or universities with responsibility or capability to fund, authorize, or carry out the corresponding recovery task.

**RECOVERY PLAN IMPLEMENTATION SCHEDULE
WHITE STURGEON: KOOTENAI RIVER POPULATION**

PRIORITY Number	TASK Number	TASK DESCRIPTION	TASK DURATION (YRS)	RESPONSIBLE PARTY	COST ESTIMATES (\$1,000)						COMMENTS
					Total Cost	FY 1999	FY 2000	FY 2001	FY 2002	FY 2003	
1	111	Conduct public review of and approve new operational guidelines for Libby Dam.	1	USACE*, MFWP, FWS, BPA, DFO, MELP	120	40	80				Begin in 1999
1	112	Implement new operational guidelines to provide annual flow regimes to benefit white sturgeon in the Kootenai River basin.	Continual	USACE*, BPA, MFWP, FWS	Unk						Costs include foregone power production; possible flood control costs, need to be determined.
1	113	Coordinate Libby Dam flow releases during April through August to achieve an optimum combination of water temperature and discharge volume.	Continual	USACE, BPA, BR, NMFS, FWS, MFWP, BC HYDRO	Unk						Instream management to fine-tune augmentation
1	114	Store water in Koocanusa Reservoir prior to spring runoff to achieve white sturgeon flow targets.	Ongoing	USACE*, BPA, MFWP	Unk						
1	115	Conduct agency coordination for implementing white sturgeon flow augmentation program.	Ongoing	USACE, BPA, BC HYDRO, DFO, MELP, MFWP, IDFG	15	3	3	3	3	3	U.S./Canada coordination.
1	116	Kootenay Lake Evaluations.	Ongoing	USACE, MELP, FQ, BPA, USFW	Unk		Unk				
1	117	Evaluate alternatives increasing peak Kootenai River flows.	Continual	DFQ, BPA, USFWS	Unk			Unk			
1	118	Use existing authorities to conserve and restore Kootenai River white sturgeon.	Continual	USACE, FWS, BPA, NMFS	Unk						Section 7 consultation on Libby Dam operations.
1	121	Monitor potential residential or agricultural flooding, levee erosion, and groundwater seepage resulting from flow augmentation.	Ongoing	USACE	Unk						Concurrent with Task 112.
1	122	Identify opportunities to restore natural floodplain functions along the Kootenai River.	Ongoing	USACE*, NRCS, DFO, MELP	Unk						
1	123	Develop public information program to explain USACE past and current flood control compensation program.	1-2	USACE	Unk	Unk	Unk				1 - 2 year public information program.
1	124	Monitor impacts of flow augmentation on Kootenai River levees and Kootenay Lake in British Columbia.	Continual	DFO, MELP	Unk						New monitoring program may be needed.
1	125	Assess consolidation of white sturgeon spawning and incubation, habitat quality, and potential substrate improvement measures.	1-2	FWS, IDFG, KTOI	***						Funded as part of Task 321

* - Lead Agency
 *** - Costs associated as part of other recovery tasks.

unk - Cost estimates are unknown.
 ongoing - Task is currently being implemented.
 continual - Task will be implemented annually when approved and/or funded.

**RECOVERY PLAN IMPLEMENTATION SCHEDULE
WHITE STURGEON: KOOTENAI RIVER POPULATION**

PRIORITY Number	TASK Number	TASK DESCRIPTION	TASK DURATION (YRS)	RESPONSIBLE PARTY	COST ESTIMATES (\$1,000)					COMMENTS	
					Total Cost	FY 1999	FY 2000	FY 2001	FY 2002		FY 2003
1	211	Obtain necessary local, State, Tribal, Federal, and Canadian approval and permits for all conservation aquaculture activities.	Continual	IDFG, KTOI, FWS, MFWP	Unk						Need to obtain permits annually e.g. Section 10.
1	221	Determine water quality standards for KTOI Hatchery.	1	BPA, KTOI* MFWP	***						1 year, as part of Task 224
1	222	Upgrade KTOI hatchery to meet conservation aquaculture objectives.	2 - 3	BPA, KTOI,	1711						Complete upgrade begun in 1998. Cost accrued beginning in 1998.
1	223	Maintain Kootenai Trout trout hatchery as a secondary rearing facility.	1	BPA, MFWP	310	61	62	63	62	62	Contract costs part of dollars allocated as part of task 222
1	224	Implement the conservation aquaculture program.	10	BPA, KTOI,	1300	240	290	260	270	280	Costs of operating existing KTOI hatchery, out year costs be higher if hatchery is expanded
1	231	Use adopted white sturgeon broodstock collection protocol.	10	BPA, KTOI,* IDFG	***						Funded as part of Task 224
1	232	Collect adequate numbers of male and female broodstock to maintain the genetic quality.	10	BPA, KTOI, IDFG	***						Funded as part of Task 224.
1	233	Annually evaluate the conservation aquaculture program.	10	BPA, KTOI IDFG, MFWP	***						Funded as part of Task 224 will be evaluated annually.
1	241	Evaluate appropriate production goals	Continual	KTOI, BPA, FWS, MFWP, IDFG, DFO, MELP	***						Funded as part of Task 224
1	242	Develop a fish health plan for hatchery	Continual	KTOI*, ALL AGENCIES	***						Funded as part of Task 224
1	243	Develop tagging protocols for hatchery reared white sturgeon.	2	KTOI, IDFG*, DFO, MELP, MFWP	Unk	Unk	Unk				Funded as part of Task 224
1	245	Evaluate feasibility of establishing an experimental white sturgeon population outside of the current occupied range.	2	ALL AGENCIES	***						Recovery team will consult with State and Canada agencies.

**RECOVERY PLAN IMPLEMENTATION SCHEDULE
WHITE STURGEON: KOOTENAI RIVER POPULATION**

PRIORITY Number	TASK Number	TASK DESCRIPTION	TASK DURATION (YRS)	RESPONSIBLE PARTY	COST ESTIMATES (\$1,000)						COMMENTS
					Total Cost	FY 1999	FY 2000	FY 2001	FY 2002	FY 2003	
1	25	Release hatchery reared white sturgeon into the Kootenai River basin.	10	KTOI*, ALL AGENCIES	Unk						Funded as part of Task 224
1	251	Adjust white sturgeon releases as necessary, to meet objectives of the Kincaid breeding plan.	10	KTOI, FWS, MFWP, BPA	***						Funded as part of Task 224
1	261	Determine factors limiting production (natural and hatchery) and habitat use patterns for each life history stage.	Ongoing	BPA, KTOI, MFWP, IDFG	Unk						Funded as part of Task 321
1	321	Describe response of spawning white sturgeon to various Kootenai River flows, water temperatures, and Kootenay Lake elevations.	Continual	KTOI, MELP MFWP, IDFG, NMFS	4000	750	775	800	825	850	120 K of this total is for monitoring effects of augmentation in Kootenai Reservoir
1	322	Measure white sturgeon spawning annually.	Continual	IDFG, KTOI, MFWP	Unk						Funded as part of Task 321
1	323	Measure white sturgeon larval, fry, and juvenile abundance and distribution in the Kootenai River and Kootenay Lake annually.	Continual	BPA, IDFG, KTOI, MFWP	Unk						Funded as part of Task 321
1	41	Develop a Participation Plan to support implementation of the Kootenai River white sturgeon recovery plan.	1	ALL AGENCIES	Unk		Unk				Will be completed as part of final recovery plan.

**RECOVERY PLAN IMPLEMENTATION SCHEDULE
WHITE STURGEON: KOOTENAI RIVER POPULATION**

PRIORITY Number	TASK Number	TASK DESCRIPTION	TASK DURATION (YRS)	RESPONSIBLE PARTY	COST ESTIMATES (\$1,000)					COMMENTS	
					Total Cost	FY 1999	FY 2000	FY 2001	FY 2002		FY 2003
2	244	Develop a policy for hatchery white sturgeon produced in excess of beneficial uses identified in this plan.	1	BPA, FWS, KTOI, IDFG, MFWP, DFO, MELP	***	Unk					May require need for Section 10 permits. Funded as part of Task 223.
2	311	Sample adult and juvenile white sturgeon in the Kootenai River and Kootenay Lake.	Ongoing	BPA, KTOI, MFWP, IDFG, BC, MELP	***						Funded as part of Task 321.
2	312	Collect and preserve tissue and blood samples for genetic analysis system.	Ongoing	BC, MELP, BPA, KTOI, MFWP, IDFG	***						Funded as part of Task 321.
2	313	Compile catch data to refine white sturgeon population size estimates annually in the Kootenai River basin.	Ongoing	BPA	***						Funded as part of Task 321.
2	314	Develop a juvenile white sturgeon year class index.	2	BPA, IDFG*	***	Unk	Unk				Funded as part of Task 321.
2	323	Measure white sturgeon larval, fry, and juvenile abundance and distribution in the Kootenai River and Kootenay Lake annually.	Ongoing	BPA	***						Funded as part of Task 321.
2	324	Quantify sturgeon spawning incubation habitat and early rearing habitat using IFIM.	2	BPA, IDFG, KTOI, MFWP	***		Unk	Unk			Begin in 1997, funded as part of Task 321.
2	331	Assess the necessity of increasing nutrients in the Kootenai River.	5	BPA, KTOI	***						Funded as part of task 321.
2	332	Use computer modeling to refine and analyze recovery tasks	1	USFWS, MELP, IDFG, KTOI	Unk	Unk					Either 1996 or 1997.
2	341	Compile existing information on contaminants in the Kootenai River.	2	BPA, KTOI* IDFG	Unk	Unk	Unk				Funded as part of Task 331.
2	342	Conduct contaminants bioassays to evaluate the effects of selected chemicals on white sturgeon.	5	BPA, KTOI*	Unk						Part of recovery Task 331
2	421	Recommend additional research and management regarding ecosystem and fishery improvement measures for the Kootenai River Basin.	Ongoing	Recovery Team	Unk						Ongoing as needed.

**RECOVERY PLAN IMPLEMENTATION SCHEDULE
WHITE STURGEON: KOOTENAI RIVER POPULATION**

PRIORITY Number	TASK Number	TASK DESCRIPTION	TASK DURATION (YRS)	RESPONSIBLE PARTY	COST ESTIMATES (\$1,000)						COMMENTS
					Total	FY 1999	FY 2000	FY 2001	FY 2002	FY 2003	
3	44	Determine the indirect recovery needs of foregone power generation	Ongoing	BPA, USACE, BCHYDRO	Unk						
n/a	511	Collect baseline abundance, distribution, and reproduction date for kokanee in Idaho Kootenai River tributaries.	5	IDFG, KTOI MFWP	809	147	155	162	169	176	Ongoing
n/a	521	Determine distribution and life history of burbot.	5	IDFG*, MFWP	735	133	140	147	154	161	Ongoing
n/a	522	Determine secondary impacts from proposed flow augmentation program on burbot reproduction.	2	IDFG*, MFWP	Unk	Unk					Costs part of Task 521.
n/a	531	Determine the status, distribution, and habitat use of rainbow trout in the Kootenai River from Libby Dam to Bonners Ferry.	5	IDFG, MFWP	900	170	175	180	185	190	Ongoing
n/a	541	Determine distribution and status of bull trout in in tributaries of the Kootenai River.	2	IDFG*, MFWP	300	54	57	60	63	66	Started in 1996, ongoing
n/a	542	Identify additional conservation measures to protect bull trout in the Kootenai River drainage in Montana, Idaho, and B.C.	1	BPA, MFWP IDFG	Unk			Unk			Using results from Task 541.
n/a	551	Determine the secondary effects of proposed white sturgeon flow augmentation on mountain whitefish.	4	BPA, MFWP	75	13	14	15	16	17	
3	561	Effects of flow augmentation on TGP in the Kootenay River and Columbia River downstream of Kootenay Lake.	?	DFO, MELP	Unk						Canada projects
3	562	Effects of flow augmentation on the Duncan and Arrow Reservoir/Lower systems	?	DFO, MELP	Unk						Canada projects.
3	6	Assess the overall success of implementation of the recovery plan and revise accordingly.	Ongoing	Recovery Team	Unk						