

Appendix 2. Summary of coho salmon carcasses recovered during the 1989-90 mainstem Trinity River spawner survey.

Survey week	Date begun	Ad-clips a/	Program marks b/	Males	Female coho			Percent unspawned	Week totals
					unknown spawned c/	Spawned	Unspawned		
7	30-Oct	0	0	1	0	0	1	100.0	2
8	06-Nov	0	2		5	0	2	100.0	10
9	13-Nov	0	3	7	14	5	1	16.7	30
10	20-Nov	1	2	19	11	16	4	20.0	53
11	27-Nov	2	9	43	7	98	6	5.8	165
12	04-Dec	0	24	129	7	115	5	4.2	280
13	11-Dec	0	15	98	0	100	9	8.3	222
14	18-Dec	0	6	132	0	98	5	4.9	241
15	25-Dec	0	7	54	0	84	4	4.5	149
16	01-Jan	0	5	31	0	74	3	3.9	113
17	05-Jan	0	2	22	0	42	1	2.3	67
18	15-Jan	0	0	9	0	11	2	15.4	22
19	22-Jan	0	0	3	0	3	0	0.0	6
Totals:		3	75	549	44	646	43		1,360
Average:								6.2	

a/ Adipose fin-clipped fish.

b/ Includes coho salmon which were previously marked (spaghetti tag/operculum punch) at various sites downstream of the survey area.

c1 Includes female for which spawning condition was not assessed.