

**TABLE 7.** Program mark recoveries from condition-one chinook salmon during the 1990-91 mainstem Trinity River spawner survey.

| Tag site                          | spring-run chinook          |                              |                 | Fall-run chinook |                |                 |
|-----------------------------------|-----------------------------|------------------------------|-----------------|------------------|----------------|-----------------|
|                                   | Program marks <sup>a/</sup> | Total observed <sup>b/</sup> | % Program marks | program marks    | Total observed | % Program marks |
| Willow Creek Weir <sup>c/</sup>   | 0                           | 295                          | 0.0             | 10               | 238            | 4.2             |
| Junction City Weir                | 28                          | 295                          | 9.5             | 25 <sup>d/</sup> | 238            | 10.5            |
| Klamath River mouth <sup>c/</sup> | 0                           | 295                          | 0.0             | 1                | 238            | 0.4             |
| <b>Totals:</b>                    | <b>28</b>                   |                              |                 | <b>36</b>        |                |                 |

<sup>a/</sup> Program marks include spaghetti tags and operculum punches.

<sup>b/</sup> Total number of condition-one chinook salmon observed during the mainstem Trinity River spawner survey.

<sup>c/</sup> Only fall-run chinook salmon were tagged at these sites.

<sup>d/</sup> Includes two Program marks which were also observed at Willow Creek Weir.