

Appendix 4. Summary of coho salmon carcasses recovered during the 1991-92 mainstem Trinity River spawner survey.

Survey week	Date begun	Ad-clips a/	Program marks b/	Males	Female coho			Week totals	
					Spawned	Unspawned	Percent unspawned Unknown c/		
1	16-Sep	0	0	0	0	0	—	0	0
2	23-Sep	0	0	0	0	0	—	0	0
3	30-Sep	0	0	0	0	0	—	0	0
4	07-Oct	0	0	0	0	0	—	0	0
5	14-Oct	0	0	0	0	0	—	0	0
6	21-Oct	0	0	1	1	0	0	0	2
7	28-Oct	0	0	1	5	0	0	0	6
8	04-Nov	0	1	1	4	0	0	0	6
9	11-Nov	0	5	4	9	0	0	1	19
10	18-Nov	0	5	11	14	0	0	0	30
11	25-Nov	1	1	12	9	0	0	0	23
12	02-Dec	0	3	11	14	0	0	0	28
13	09-Dec	0	1	1	8	0	0	0	10
14	16-Dec	0	1	2	0	0	—	0	3
Totals:		1	17	44	64	0		1	127
Average:							0		

a/ Adipose fin-clipped coho salmon.

b/ Includes coho salmon which were previously marked (spaghetti tags/operculum punched) downstream of the survey area.

c/ Includes female coho for which spawning condition was not assessed.